

thm_2Ebool_2ESELECT__REFL (TMNYYB-
wXXV7XY69wz249wn4EZFhnHz8pVc2)

October 26, 2020

Definition 1 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (ap P x)) \text{ then } (the (\lambda x. x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define `c_2Ebool_2ET` to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V 0x \in 2. V 0x)) (\lambda V 1x \in 2. V 1x))$

Definition 4 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V 0P \in (2^{A-27a}). (ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V 1y \in A. 27a. (ap (ap (c_2Emin_2E_40 A. 27a) (\lambda V 1y \in A. 27a. (ap (ap (c_2Emin_2E_3D A. 27a) V 1y) V 0x))) = V 0x))$

Theorem 1

$$\forall A. 27a. nonempty A. 27a \Rightarrow (\forall V 0x \in A. 27a. ((ap (c_2Emin_2E_40 A. 27a) (\lambda V 1y \in A. 27a. (ap (ap (c_2Emin_2E_3D A. 27a) V 1y) V 0x))) = V 0x))$$