

thm_2Ebool_2EU EXISTS_SIMP (TMVqNHiwA8aLvrzKEAQj9ixCtnyS9a4irbE)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow P \Rightarrow Q)$ of type ι .

Definition 2 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define `c_2Ebool_2ET` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

Definition 4 We define `c_2Ebool_2E_21` to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A_27a}))$

Definition 5 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V2t \in 2. V2t)))$

Definition 6 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (the (\lambda x. x \in A \wedge p (\text{ap } P x)))$ of type $\iota \Rightarrow \iota$.

Definition 7 We define `c_2Ebool_2E_3F` to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (\text{ap } V0P (\text{ap } (\text{c_2Emin_2E_40 } A_27a P))))$

Definition 8 We define `c_2Ebool_2E_3F_21` to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (\text{ap } (\text{ap } (\text{c_2Ebool_2E_2F_5C } A_27a V0P))))$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p \Rightarrow V0t1) \Rightarrow (p \Rightarrow V1t2)) \Rightarrow (((p \Rightarrow V1t2) \Rightarrow (p \Rightarrow V0t1)) \Rightarrow ((p \Rightarrow V0t1) \Leftrightarrow (p \Rightarrow V1t2)))))) \quad (1)$$

Assume the following.

$$\forall A_27a. \text{nonempty } A_27a \Rightarrow (\forall V0t \in 2. ((\exists V1x \in A_27a. (p \Rightarrow V0t)) \Leftrightarrow (p \Rightarrow V0t))) \quad (2)$$

Assume the following.

$$\forall A_27a. \text{nonempty } A_27a \Rightarrow (\forall V0P \in 2. (\forall V1Q \in (2^{A_27a}). ((\forall V2x \in A_27a. ((p \Rightarrow V0P) \Rightarrow (p (\text{ap } V1Q V2x)))) \Leftrightarrow ((p \Rightarrow V0P) \Rightarrow (\forall V3x \in A_27a. (p (\text{ap } V1Q V3x))))))) \quad (3)$$

Theorem 1

$$\forall A_27a. \text{nonempty } A_27a \Rightarrow (\forall V0t \in 2. ((p (\text{ap } (\text{c_2Ebool_2E_3F_21 } A_27a) (\lambda V1x \in A_27a. V0t))) \Leftrightarrow ((p \Rightarrow V0t) \wedge (\forall V2x \in A_27a. (\forall V3y \in A_27a. (V2x = V3y))))))$$