

# thm\_2Ebool\_2Ebool\_\_case\_\_CONG (TM- FwXK95kuEGj2ZMi2KbDUTMrzGPfrzsFAS)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D` to be  $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj\_o } (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define `c_2Ebool_2ET` to be  $(\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

**Definition 3** We define `c_2Ebool_2E_21` to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^{A\_27a}))$

**Definition 4** We define `c_2Ebool_2EF` to be  $(\text{ap } (\text{c\_2Ebool\_2E\_21 } 2) (\lambda V0t \in 2. V0t))$ .

**Definition 5** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2. \lambda Q \in 2. \text{inj\_o } (p \Rightarrow Q)$  of type  $\iota$ .

**Definition 6** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2) (\lambda V2t \in 2. V2t))$

**Definition 7** We define `c_2Emin_2E_40` to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 8** We define `c_2Ebool_2ECOND` to be  $\lambda A\_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A\_27a. (\lambda V2t2 \in A\_27a. (\text{ap } (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D\_3D\_3E } V0t) (\text{c\_2Ebool\_2E\_21 } V1t1) V2t2))$

**Definition 9** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D\_3D\_3E } V0t) (\text{c\_2Ebool\_2E\_21 } V0t) V0t))$

Assume the following.

$$\begin{aligned}
 & \forall A\_27a. \text{nonempty } A\_27a \Rightarrow (\forall V0P \in 2. (\forall V1Q \in 2. \\
 & (\forall V2x \in A\_27a. (\forall V3x\_27 \in A\_27a. (\forall V4y \in A\_27a. \\
 & (\forall V5y\_27 \in A\_27a. (((p \Rightarrow V0P) \Leftrightarrow (p \Rightarrow V1Q)) \wedge ((p \Rightarrow V1Q) \Rightarrow (V2x = V3x\_27)) \wedge \\
 & ((\neg(p \Rightarrow V1Q)) \Rightarrow (V4y = V5y\_27)))))) \Rightarrow ((\text{ap } (\text{ap } (\text{ap } (\text{c\_2Ebool\_2ECOND } A\_27a) \\
 & V0P) V2x) V4y) = (\text{ap } (\text{ap } (\text{ap } (\text{c\_2Ebool\_2ECOND } A\_27a) V1Q) V3x\_27) \\
 & V5y\_27)))))))))
 \end{aligned} \tag{1}$$

**Theorem 1**

$$\begin{aligned}
 & \forall A\_27a. \text{nonempty } A\_27a \Rightarrow (\forall V0P \in 2. (\forall V1Q \in 2. \\
 & (\forall V2x \in A\_27a. (\forall V3x\_27 \in A\_27a. (\forall V4y \in A\_27a. \\
 & (\forall V5y\_27 \in A\_27a. (((p \Rightarrow V0P) \Leftrightarrow (p \Rightarrow V1Q)) \wedge ((p \Rightarrow V1Q) \Rightarrow (V2x = V3x\_27)) \wedge \\
 & ((\neg(p \Rightarrow V1Q)) \Rightarrow (V4y = V5y\_27)))))) \Rightarrow ((\text{ap } (\text{ap } (\text{ap } (\text{c\_2Ebool\_2ECOND } A\_27a) \\
 & V0P) V2x) V4y) = (\text{ap } (\text{ap } (\text{ap } (\text{c\_2Ebool\_2ECOND } A\_27a) V1Q) V3x\_27) \\
 & V5y\_27)))))))))
 \end{aligned}$$