

# thm\_2Ebool\_2Eitself\_Axiom (TMRo- QipFq1ezqMDf3WBo8MR9wsq4HiVUYhC)

October 26, 2020

Let  $ty\_2Ebool\_2Eitself : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty\_2Ebool\_2Eitself\ A0) \quad (1)$$

Let  $c\_2Ebool\_2Ethe\_value : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Ebool\_2Ethe\_value\ A\_27a \in (ty\_2Ebool\_2Eitself\ A\_27a) \quad (2)$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o\ (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if\ (\exists x \in A.p\ (ap\ P\ x))\ \mathbf{then}\ (the\ (\lambda x.x \in A \wedge p\ x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40\ A\_27a)\ V0P)))$

**Definition 4** We define  $c\_2Ebool\_2E\_T$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x)))$

**Definition 5** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A\_27a})\ V0P))))$

**Theorem 1**

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow (\forall V0e \in A\_27b.(\exists V1f \in (A\_27b^{(ty\_2Ebool\_2Eitself\ A\_27a)}).((ap\ V1f\ (c\_2Ebool\_2Ethe\_value\ A\_27a)) = V0e)))$$