

thm_2Ecardinal_2ECOUNTABLE__ALT__cardleq
(TMWnx78maQJo5aaphWQBxr68qcv14opN8Kv)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 4 We define $c_2Ebool_2E_2IN$ to be $\lambda A.\lambda a : \iota.(\lambda V0x \in A.\lambda a.\lambda V1f \in (2^{A-27a}).(ap V1f V0x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 7 We define $c_2Epred_set_2EINJ$ to be $\lambda A.\lambda a : \iota.\lambda A.\lambda b : \iota.\lambda V0f \in (A \rightarrow 2^{A-27a}).\lambda V1s \in (2^{A-27a})$

Definition 8 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$. **if** $(\exists x \in A.p (ap P x))$ **then** $(the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 9 We define $c_2Ebool_2E_3F$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40 A P))))$

Definition 10 We define $c_2Ecardinal_2Ecardleq$ to be $\lambda A.\lambda a : \iota.\lambda A.\lambda b : \iota.\lambda V0s1 \in (2^{A-27a}).\lambda V1s2 \in (2^{A-27a})$

Definition 11 We define $c_2Epred_set_2EUNIV$ to be $\lambda A.\lambda a : \iota.(\lambda V0x \in A.\lambda a.c_2Ebool_2E_2T)$.

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Definition 12 We define $c_2Ecardinal_2Ecardgeq$ to be $\lambda A.\lambda a : \iota.\lambda A.\lambda b : \iota.\lambda V0s \in (2^{A-27a}).\lambda V1t \in (2^{A-27a})$

Definition 13 We define $c_2Epred_set_2Ecountable$ to be $\lambda A.\lambda a : \iota.\lambda V0s \in (2^{A-27a}).(ap (c_2Ebool_2E_3F A s))$

Assume the following.

$$True \quad (2)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (3)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (4)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0s \in (2^{A_27a}). (\forall V1t \in (2^{A_27b}). ((p\ (ap\ (ap\ (c_2Ecardinal_2Ecardgeq\ A_27a\ A_27b)\ V0s)\ V1t)) \Leftrightarrow (p\ (ap\ (ap\ (c_2Ecardinal_2Ecardleq\ A_27b\ A_27a)\ V1t)\ V0s)))))) \quad (5)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in (2^{A_27a}). ((p\ (ap\ (c_2Epred_set_2Ecountable\ A_27a)\ V0t)) \Leftrightarrow (p\ (ap\ (ap\ (c_2Ecardinal_2Ecardgeq\ ty_2Enum_2Enum\ A_27a)\ (c_2Epred_set_2EUNIV\ ty_2Enum_2Enum))\ V0t)))))) \quad (6)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0s \in (2^{A_27a}). ((p\ (ap\ (c_2Epred_set_2Ecountable\ A_27a)\ V0s)) \Leftrightarrow (p\ (ap\ (ap\ (c_2Ecardinal_2Ecardleq\ A_27a\ ty_2Enum_2Enum)\ V0s)\ (c_2Epred_set_2EUNIV\ ty_2Enum_2Enum))))))$$