

thm\_2Ecardinal\_2ECOUNTABLE\_\_RESTRICT  
(TMRrjrL-  
DAUEm1UUtAVDERuEahHsVVrjJA86)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Epair\_2Eprod A0 A1) \tag{1}$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epair\_2EABS\_prod A\_27a A\_27b \in ((ty\_2Epair\_2Eprod A\_27a A\_27b)^{(2^{A\_27b})^{A\_27a}}) \tag{2}$$

**Definition 8** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2E$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40$

Let  $c\_2Epred\_set\_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epred\_set\_2EGSPEC \\ A\_27a\ A\_27b \in ((2^{A\_27a})^{(ty\_2Epair\_2Eprod\ A\_27a\ 2)^{A\_27b}}) \end{aligned} \quad (3)$$

**Definition 11** We define  $c\_2Ebool\_2EIN$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. (\lambda V1f \in (2^{A\_27a}). (ap\ V1f\ V0x)))$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (4)$$

**Definition 12** We define  $c\_2Epred\_set\_2EUNIV$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. c\_2Ebool\_2EIN)$ .

**Definition 13** We define  $c\_2Epred\_set\_2EINJ$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0f \in (A\_27b^{A\_27a}). \lambda V1s \in (2^{A\_27a})$

**Definition 14** We define  $c\_2Epred\_set\_2Ecountable$  to be  $\lambda A\_27a : \iota. \lambda V0s \in (2^{A\_27a}). (ap\ (c\_2Ebool\_2EIN\ 3F$

**Definition 15** We define  $c\_2Epred\_set\_2ESUBSET$  to be  $\lambda A\_27a : \iota. \lambda V0s \in (2^{A\_27a}). \lambda V1t \in (2^{A\_27a}). (ap\ ($

Assume the following.

$$True \quad (5)$$

Assume the following.

$$\begin{aligned} (\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p \\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in \\ A\_27a. (p\ V0t) \Leftrightarrow (p\ V0t))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (( \\ (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in \\ A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg (p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg ( \\ p\ V0t)))))) \end{aligned} \quad (10)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow (p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (11)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x_{.27} \in 2.(\forall V2y \in 2.(\forall V3y_{.27} \in 2.(((p V0x) \Leftrightarrow (p V1x_{.27})) \wedge ((p V1x_{.27}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{.27})))))) \Rightarrow (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{.27}) \Rightarrow (p V3y_{.27})))))) \quad (12)$$

Assume the following.

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0P \in (2^{A_{.27a}}).(\forall V1a \in A_{.27a}.((\exists V2x \in A_{.27a}.((V2x = V1a) \wedge (p (ap V0P V2x)))) \Leftrightarrow (p (ap V0P V1a)))))) \quad (13)$$

Assume the following.

$$(\forall V0r \in 2.(\forall V1p \in 2.(\forall V2q \in 2.(((p V1p) \wedge (p V2q) \Rightarrow (p V0r)) \Leftrightarrow ((p V1p) \Rightarrow ((p V2q) \Rightarrow (p V0r)))))) \quad (14)$$

Assume the following.

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow \forall A_{.27b}.nonempty A_{.27b} \Rightarrow (\forall V0x \in A_{.27a}.(\forall V1y \in A_{.27b}.(\forall V2a \in A_{.27a}.(\forall V3b \in A_{.27b}.(((ap (ap (c_{.2E}pair_{.2E} C A_{.27a} A_{.27b}) V0x) V1y) = (ap (ap (c_{.2E}pair_{.2E} C A_{.27a} A_{.27b}) V2a) V3b)) \Leftrightarrow ((V0x = V2a) \wedge (V1y = V3b)))))) \quad (15)$$

Assume the following.

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow \forall A_{.27b}.nonempty A_{.27b} \Rightarrow (\forall V0f \in ((ty_{.2E}pair_{.2E}prod A_{.27a} 2)^{A_{.27b}}).(\forall V1v \in A_{.27a}.((p (ap (ap (c_{.2E}bool_{.2E} IN A_{.27a}) V1v) (ap (c_{.2E}pred_{.set} EGSPEC A_{.27a} A_{.27b}) V0f))) \Leftrightarrow (\exists V2x \in A_{.27b}.((ap (ap (c_{.2E}pair_{.2E} C A_{.27a} 2) V1v) c_{.2E}bool_{.2E} ET) = (ap V0f V2x)))))) \quad (16)$$

Assume the following.

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0s \in (2^{A_{.27a}}).(\forall V1t \in (2^{A_{.27a}}).(((p (ap (ap (c_{.2E}pred_{.set} E SUBSET A_{.27a}) V0s) V1t)) \wedge (p (ap (c_{.2E}pred_{.set} E countable A_{.27a}) V1t))) \Rightarrow (p (ap (c_{.2E}pred_{.set} E countable A_{.27a}) V0s)))))) \quad (17)$$

### Theorem 1

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0s \in (2^{A_{.27a}}).(\forall V1P \in (2^{A_{.27a}}).((p (ap (c_{.2E}pred_{.set} E countable A_{.27a}) V0s)) \Rightarrow (p (ap (c_{.2E}pred_{.set} E countable A_{.27a}) (ap (c_{.2E}pred_{.set} EGSPEC A_{.27a} A_{.27a}) (\lambda V2x \in A_{.27a}.(ap (ap (c_{.2E}pair_{.2E} C A_{.27a} 2) V2x) (ap (ap (c_{.2E}bool_{.2E} F_{.5C} (ap (ap (c_{.2E}bool_{.2E} IN A_{.27a}) V2x) V0s)) (ap V1P V2x))))))))))$$