

thm_2Ecardinal_2EEMPTY_CARDLEQ (TMLfoH1kHzzBpwHdnThxa26QobApjgSXv5U)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2T` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2))) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x)$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 4 We define `c_2Ebool_2E_2F` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V0t \in 2. V0t)$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t)) \text{c_2Ebool_2E_2F}))$

Definition 7 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 8 We define `c_2Ebool_2E_3F` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } V0P (\text{ap } (\text{c_2Emin_2E_40 } A))))$

Definition 9 We define `c_2Ebool_2E_2IN` to be $\lambda A. 27a : \iota. (\lambda V0x \in A. 27a. (\lambda V1f \in (2^{A-27a}). (\text{ap } V1f V0x)))$

Definition 10 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

Definition 11 We define `c_2Epred_set_2EINJ` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0f \in (A. 27b^{A-27a}). \lambda V1s \in (2^{A-27a}).$

Definition 12 We define `c_2Ecardinal_2Ecardleq` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0s1 \in (2^{A-27a}). \lambda V1s2 \in (2^{A-27a}).$

Definition 13 We define `c_2Epred_set_2EEMPTY` to be $\lambda A. 27a : \iota. (\lambda V0x \in A. 27a. \text{c_2Ebool_2E_2F}).$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow (\forall V0t \in 2. ((\exists V1x \in A. 27a. (p V0t)) \Leftrightarrow (p V0t))) \tag{2}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow True) \Leftrightarrow \\
& (p \ V0t)) \wedge (((False \Leftrightarrow (p \ V0t)) \Leftrightarrow \neg(p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\
& p \ V0t))))))
\end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty \ A_27a \Rightarrow \forall A_27b.nonempty \ A_27b \Rightarrow (\\
& \quad \forall V0f \in (A_27b^{A_27a}).((\forall V1s \in (2^{A_27b}).(p \ (ap \ (ap \\
& (ap \ (c_2Epred_set_2EINJ \ A_27a \ A_27b) \ V0f) \ (c_2Epred_set_2EEMPTY \\
& A_27a)) \ V1s))) \wedge (\forall V2s \in (2^{A_27a}).((p \ (ap \ (ap \ (ap \ (c_2Epred_set_2EINJ \\
& A_27a \ A_27b) \ V0f) \ V2s) \ (c_2Epred_set_2EEMPTY \ A_27b))) \Leftrightarrow (V2s = \\
& (c_2Epred_set_2EEMPTY \ A_27a))))))
\end{aligned} \tag{4}$$

Theorem 1

$$\begin{aligned}
& \forall A_27a.nonempty \ A_27a \Rightarrow \forall A_27b.nonempty \ A_27b \Rightarrow (\\
& \quad \forall V0t \in (2^{A_27b}).(p \ (ap \ (ap \ (c_2Ecardinal_2Ecardleq \ A_27a \\
& A_27b) \ (c_2Epred_set_2EEMPTY \ A_27a)) \ V0t)))
\end{aligned}$$