

thm\_2Ecardinal\_2EGE  
 (TMaCS3SwAuMHA7Yf6qjKVSwo77VhkKhnaoV)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (1)$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})) (\lambda V1P \in 2.V1P)))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2. \lambda Q \in 2. inj\_o (p \ P \Rightarrow p \ Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2. (ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2. inj\_o (V0t1 = V1t2))))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (2)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (3)$$

Let  $c\_2Enum\_2EAABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EAABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (4)$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. (ap c\_2Enum\_2EAABS\_num m)$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (ap P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 12** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 13** We define  $c\_2Earithmetic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 14** We define  $c\_2Earithmetic\_2E\_3E$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 15** We define  $c\_2Earithmetic\_2E\_3E\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum.(\forall V1m \in ty\_2Enum\_2Enum.((p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3E\_3D\ V0n)\ V1m)) \Leftrightarrow (p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3C\_3D\ V1m)\ V0n)))))) \quad (5)$$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2))))) \quad (7)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (8)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True)))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (10)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (11)$$

Assume the following.

$$(\forall V0A \in 2.((p\ V0A) \Rightarrow ((\neg(p\ V0A)) \Rightarrow False))) \quad (12)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((p V0A) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow False) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (13)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (14)$$

Assume the following.

$$(\forall V0A \in 2. (((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (15)$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p V0p) \Leftrightarrow (p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee ((\neg(p V1q) \vee (p V2r)) \vee (\neg(p V1q)) \vee ((p V2r) \vee ((\neg(p V1q) \vee (\neg(p V0p))))))))))) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p V0p) \Leftrightarrow (p V1q) \wedge (p V2r))) \Leftrightarrow (((p V0p) \vee ((\neg(p V1q) \vee (\neg(p V2r)))) \wedge (((p V1q) \vee (\neg(p V0p)) \wedge ((p V2r) \vee (\neg(p V0p))))))))))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p V0p) \Leftrightarrow (p V1q) \vee (p V2r))) \Leftrightarrow (((p V0p) \vee (\neg(p V1q))) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge (((p V1q) \vee ((p V2r) \vee (\neg(p V0p))))))))))) \end{aligned} \quad (18)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (((p V0p) \Leftrightarrow (\neg(p V1q))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge ((\neg(p V1q)) \vee (\neg(p V0p))))))) \quad (19)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p)))) \quad (20)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (\neg(p V1q)))))) \quad (21)$$

### Theorem 1

$$\begin{aligned} & (\forall V0n \in ty\_2Enum\_2Enum. (\forall V1m \in ty\_2Enum\_2Enum. ((p (ap (ap c_2Earithmetic_2E_3E_3D V1m) V0n)) \Leftrightarrow (p (ap (ap c_2Earithmetic_2E_3C_3D V0n) V1m))))) \end{aligned}$$