

thm_2Ecardinal_2ESURJECTIVE__ON__RIGHT__INVERSE (TMQ7BN1Y4bRBWPpN7tWxP3Y41dbuqVeiXeJ)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 6 We define $c_2Ebool_2E_2IN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 7 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 8 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a P))))$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \tag{2}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \tag{3}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{4}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in 2. (\forall V1Q \in (2^{A_27a}). ((\exists V2x \in A_27a. ((p\ V0P) \Rightarrow (p\ (ap\ V1Q\ V2x)))) \Leftrightarrow ((p\ V0P) \Rightarrow (\exists V3x \in A_27a. (p\ (ap\ V1Q\ V3x)))))))) \quad (5)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0P \in ((2^{A_27b})^{A_27a}). ((\forall V1x \in A_27a. (\exists V2y \in A_27b. (p\ (ap\ (ap\ V0P\ V1x)\ V2y)))) \Leftrightarrow (\exists V3f \in (A_27b^{A_27a}). (\forall V4x \in A_27a. (p\ (ap\ (ap\ V0P\ V4x)\ (ap\ V3f\ V4x)))))))) \quad (6)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0s \in (2^{A_27a}). (\forall V1f \in (A_27b^{A_27a}). (\forall V2t \in (2^{A_27b}). ((\forall V3y \in A_27b. ((p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27b)\ V3y)\ V2t)) \Rightarrow (\exists V4x \in A_27a. ((p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27a)\ V4x)\ V0s)) \wedge ((ap\ V1f\ V4x) = V3y)))))) \Leftrightarrow (\exists V5g \in (A_27a^{A_27b}). (\forall V6y \in A_27b. ((p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27b)\ V6y)\ V2t)) \Rightarrow ((p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27a)\ (ap\ V5g\ V6y))\ V0s)) \wedge ((ap\ V1f\ (ap\ V5g\ V6y)) = V6y)))))))))) \quad (7)$$