

thm\_2Ecardinal\_2Ecardeq\_REFL  
(TMSMaPLBs5XRZ8KaXYabpME8KYEX5kgitas)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2EF$

**Definition 7** We define  $c\_2Ebool\_2E\_2EIN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$

**Definition 8** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$  **then** (the  $(\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ ).

**Definition 10** We define  $c\_2Ebool\_2E\_23F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Epred\_set\_2ESURJ$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0f \in (A\_27b^{A\_27a}).\lambda V1s \in (2^{A\_27b})$

**Definition 12** We define  $c\_2Epred\_set\_2EINJ$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0f \in (A\_27b^{A\_27a}).\lambda V1s \in (2^{A\_27b})$

**Definition 13** We define  $c\_2Epred\_set\_2EBIJ$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0f \in (A\_27b^{A\_27a}).\lambda V1s \in (2^{A\_27b})$

**Definition 14** We define  $c\_2Ecardinal\_2Ecardeq$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0s1 \in (2^{A\_27a}).\lambda V1s2 \in (2^{A\_27b})$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p V0t) \Leftrightarrow (p V0t))) \tag{2}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \wedge True) \Leftrightarrow \\
& (p \ V0t)) \wedge (((False \wedge (p \ V0t)) \Leftrightarrow False) \wedge (((p \ V0t) \wedge False) \Leftrightarrow False) \wedge \\
& (((p \ V0t) \wedge (p \ V0t)) \Leftrightarrow (p \ V0t))))))
\end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Rightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Rightarrow True) \Leftrightarrow \\
& True) \wedge (((False \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge (((p \ V0t) \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge (( \\
& (p \ V0t) \Rightarrow False) \Leftrightarrow (\neg(p \ V0t))))))
\end{aligned} \tag{4}$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \tag{5}$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{6}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow True) \Leftrightarrow \\
& (p \ V0t)) \wedge (((False \Leftrightarrow (p \ V0t)) \Leftrightarrow (\neg(p \ V0t))) \wedge (((p \ V0t) \Leftrightarrow False) \Leftrightarrow (\neg( \\
& p \ V0t))))))
\end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p \ V0t1) \Rightarrow \\
& ((p \ V1t2) \Rightarrow (p \ V2t3))) \Leftrightarrow (((p \ V0t1) \wedge (p \ V1t2)) \Rightarrow (p \ V2t3))))))
\end{aligned} \tag{8}$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in 2.(\forall V1x\_27 \in 2.(\forall V2y \in 2.(\forall V3y\_27 \in \\
& 2.(((p \ V0x) \Leftrightarrow (p \ V1x\_27)) \wedge ((p \ V1x\_27) \Rightarrow ((p \ V2y) \Leftrightarrow (p \ V3y\_27)))) \Rightarrow \\
& (((p \ V0x) \Rightarrow (p \ V2y)) \Leftrightarrow ((p \ V1x\_27) \Rightarrow (p \ V3y\_27))))))
\end{aligned} \tag{9}$$

Assume the following.

$$\begin{aligned}
& \forall A\_27a.nonempty \ A\_27a \Rightarrow \forall A\_27b.nonempty \ A\_27b \Rightarrow ( \\
& \forall V0f \in (A\_27b^{A\_27a}).(\forall V1s \in (2^{A\_27a}).(\forall V2t \in \\
& (2^{A\_27b}).((p \ (ap \ (ap \ (ap \ (c\_2Epred\_set\_2EBIJ \ A\_27a \ A\_27b) \ V0f) \\
& V1s) \ V2t)) \Leftrightarrow ((\forall V3x \in A\_27a.((p \ (ap \ (ap \ (c\_2Ebool\_2EIN \ A\_27a) \\
& V3x) \ V1s)) \Rightarrow (p \ (ap \ (ap \ (c\_2Ebool\_2EIN \ A\_27b) \ (ap \ V0f \ V3x)) \ V2t)))) \wedge \\
& (\exists V4g \in (A\_27a^{A\_27b}).((\forall V5x \in A\_27b.((p \ (ap \ (ap \ (c\_2Ebool\_2EIN \\
& A\_27b) \ V5x) \ V2t)) \Rightarrow (p \ (ap \ (ap \ (c\_2Ebool\_2EIN \ A\_27a) \ (ap \ V4g \ V5x)) \\
& V1s)))) \wedge ((\forall V6x \in A\_27a.((p \ (ap \ (ap \ (c\_2Ebool\_2EIN \ A\_27a) \\
& V6x) \ V1s)) \Rightarrow ((ap \ V4g \ (ap \ V0f \ V6x)) = V6x))) \wedge (\forall V7x \in A\_27b.( \\
& (p \ (ap \ (ap \ (c\_2Ebool\_2EIN \ A\_27b) \ V7x) \ V2t)) \Rightarrow ((ap \ V0f \ (ap \ V4g \ V7x)) = \\
& V7x))))))
\end{aligned} \tag{10}$$

**Theorem 1**

$\forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow (\forall V_0s \in (2^{A_{27a}}). (p (ap (c\_2Ecardinal\_2Ecard \ A_{27a} \ A_{27a}) \ V_0s) \ V_0s)))$