

thm_2Ecardinal_2Ecardleq__empty
(TMUvcG2g4dj9YPAc1CbJVMytaDeDxkZiamh)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2T` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V 0x \in 2. V 0x)) (\lambda V 1x \in 2. V 1x))$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V 0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a})) (\lambda V 1P \in (2^{A-27a}). V 1P)) (\lambda V 1x \in (2^{A-27a}). V 1x)))$

Definition 4 We define `c_2Ebool_2E_2F` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V 0t \in 2. V 0t))$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V 0t1 \in 2. (\lambda V 1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V 2t \in 2. V 2t))))$

Definition 7 We define `c_2Epred__set_2EEMPTY` to be $\lambda A. 27a : \iota. (\lambda V 0x \in A. 27a. \text{c_2Ebool_2E_2F})$.

Definition 8 We define `c_2Ebool_2E_2IN` to be $\lambda A. 27a : \iota. (\lambda V 0x \in A. 27a. (\lambda V 1f \in (2^{A-27a}). (\text{ap } V 1f V 0x)))$

Definition 9 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p (\text{ap } P x)))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define `c_2Ebool_2E_3F` to be $\lambda A. 27a : \iota. (\lambda V 0P \in (2^{A-27a}). (\text{ap } V 0P (\text{ap } (\text{c_2Emin_2E_40 } (2^{A-27a})) (\lambda V 1P \in (2^{A-27a}). V 1P))))$

Definition 11 We define `c_2Epred__set_2ESURJ` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V 0f \in (A. 27b^{A-27a}). \lambda V 1s \in (2^{A-27a}). \text{ap } V 1s V 0f$

Definition 12 We define `c_2Epred__set_2EINJ` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V 0f \in (A. 27b^{A-27a}). \lambda V 1s \in (2^{A-27a}). \text{ap } V 1s V 0f$

Definition 13 We define `c_2Epred__set_2EBIJ` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V 0f \in (A. 27b^{A-27a}). \lambda V 1s \in (2^{A-27a}). \text{ap } V 1s V 0f$

Definition 14 We define `c_2Ecardinal_2Ecardleq` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V 0s1 \in (2^{A-27a}). \lambda V 1s2 \in (2^{A-27a}). \text{ap } V 1s2 V 0s1$

Definition 15 We define `c_2Ebool_2E_7E` to be $(\lambda V 0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V 0t) (\text{c_2Ebool_2E_21 } 2)) (\lambda V 1t \in 2. V 1t)))$

Definition 16 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V 0t1 \in 2. (\lambda V 1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V 2t \in 2. V 2t))))$

Definition 17 We define `c_2Ecardinal_2Ecardleq` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V 0s1 \in (2^{A-27a}). \lambda V 1s2 \in (2^{A-27a}). \text{ap } V 1s2 V 0s1$

Assume the following.

$$True \quad (1)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (2)$$

Assume the following.

$$(\forall V0t \in 2. (((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \quad (3)$$

Assume the following.

$$((\forall V0t \in 2. ((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (4)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (5)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t))))) \quad (6)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow \forall A_27c.nonempty\ A_27c \Rightarrow (\forall V0x \in (2^{A_27a}). (((p\ (ap\ (ap\ (c_2Ecardinal_2Ecardeq\ A_27a\ A_27b)\ V0x)\ (c_2Epred_set_2EEMPTY\ A_27b))) \Leftrightarrow (V0x = (c_2Epred_set_2EEMPTY\ A_27a))) \wedge (((p\ (ap\ (ap\ (c_2Ecardinal_2Ecardeq\ A_27c\ A_27a)\ (c_2Epred_set_2EEMPTY\ A_27c))\ V0x)) \Leftrightarrow (V0x = (c_2Epred_set_2EEMPTY\ A_27a)))))) \quad (7)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0t \in (2^{A_27b}). (p\ (ap\ (ap\ (c_2Ecardinal_2Ecardleq\ A_27a\ A_27b)\ (c_2Epred_set_2EEMPTY\ A_27a))\ V0t))) \quad (8)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0s1 \in (2^{A_27a}). (\forall V1s2 \in (2^{A_27b}). ((p\ (ap\ (ap\ (c_2Ecardinal_2Ecardleq\ A_27a\ A_27b)\ V0s1)\ V1s2)) \Leftrightarrow ((\neg(p\ (ap\ (ap\ (c_2Ecardinal_2Ecardleq\ A_27b\ A_27a)\ V1s2)\ V0s1))) \vee (p\ (ap\ (ap\ (c_2Ecardinal_2Ecardleq\ A_27a\ A_27b)\ V0s1)\ V1s2)))))) \quad (9)$$

Theorem 1

$$\begin{aligned} & \forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow \forall A_{27b}. \text{nonempty } A_{27b} \Rightarrow (\\ & \forall V_0 x \in (2^{A_{27a}}). ((p (ap (ap (c_2Ecardinal_2Ecardleq A_{27a} \\ & A_{27b}) V_0 x) (c_2Epred_set_2EEMPTY A_{27b}))) \Leftrightarrow (V_0 x = (c_2Epred_set_2EEMPTY \\ & A_{27a})))) \end{aligned}$$