

thm_2Ecardinal_2Ecountable__thm
(TMRCFZ26F4oYaxH58misfXhe2sJ6Fm9fd1W)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 4 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 7 We define $c_2Epred_set_2EINJ$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (A_27b^{A_27a}).\lambda V1s \in (2^{A_27a})$

Definition 8 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 9 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a) P)))$

Definition 10 We define $c_2Ecardinal_2Ecardleq$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0s1 \in (2^{A_27a}).\lambda V1s2 \in (2^{A_27b})$

Definition 11 We define $c_2Epred_set_2EUNIV$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2ET)$.

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Definition 12 We define $c_2Epred_set_2Ecountable$ to be $\lambda A_27a : \iota.\lambda V0s \in (2^{A_27a}).(ap (c_2Ebool_2E_3F A_27a) s)$

Assume the following.

$$True \tag{2}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \tag{3}$$

Assume the following.

$$\forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow (\forall V0x \in A_{27a}. ((V0x = V0x) \Leftrightarrow \text{True})) \quad (4)$$

Theorem 1

$$\forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow (\forall V0s \in (2^{A_{27a}}). ((p \text{ (ap (c_2Epred_set_2Ecountable } A_{27a}) V0s)) \Leftrightarrow (p \text{ (ap (ap (c_2Ecardinal_2Ecardleq } A_{27a} \text{ ty_2Enum_2Enum) } V0s) (c_2Epred_set_2EUNIV \text{ ty_2Enum_2Enum}))))))$$