

thm\_2Ecardinal\_2Ege\_c  
(TMFPh5pLHkQUw4e9djStfrqnJUdq7YXWwf6)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D` to be  $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj\_o } (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2. \lambda Q \in 2. \text{inj\_o } (p \Rightarrow q)$  of type  $\iota$ .

**Definition 3** We define `c_2Ebool_2EIN` to be  $\lambda A_27a : \iota. (\lambda V0x \in A_27a. (\lambda V1f \in (2^{A_27a}). (\text{ap } V1f \ V0x)))$

**Definition 4** We define `c_2Ebool_2EET` to be  $(\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

**Definition 5** We define `c_2Ebool_2E_21` to be  $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^{A_27a}))))$

**Definition 6** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2) (\lambda V2t \in 2. V2t))))$

**Definition 7** We define `c_2Epred_set_2EINJ` to be  $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0f \in (A_27b^{A_27a}). \lambda V1s \in (2^{A_27a})$

**Definition 8** We define `c_2Emin_2E_40` to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P \ x)) \text{ then } (the (\lambda x. x \in A \wedge p (\text{ap } P \ x)))$  of type  $\iota \Rightarrow \iota$ .

**Definition 9** We define `c_2Ebool_2E_3F` to be  $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (\text{ap } V0P (\text{ap } (\text{c\_2Emin\_2E\_40 } A_27a))))$

**Definition 10** We define `c_2Ecardinal_2Ecardleq` to be  $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0s1 \in (2^{A_27a}). \lambda V1s2 \in (2^{A_27b})$

**Definition 11** We define `c_2Ecardinal_2Ecardgeq` to be  $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0s \in (2^{A_27a}). \lambda V1t \in (2^{A_27b})$

**Theorem 1**

$$\forall A_27a. \text{nonempty } A_27a \Rightarrow \forall A_27b. \text{nonempty } A_27b \Rightarrow (\forall V0s \in (2^{A_27a}). (\forall V1t \in (2^{A_27b}). ((p (\text{ap } (\text{ap } (\text{c\_2Ecardinal\_2Ecardgeq } A_27a \ A_27b) \ V0s) \ V1t)) \Leftrightarrow (p (\text{ap } (\text{ap } (\text{c\_2Ecardinal\_2Ecardleq } A_27b \ A_27a) \ V1t) \ V0s))))))$$