

thm_2Ecombin_2EAPPLY__UPDATE__ID
(TMKJQmmNv1up3fqCeAgRg6Kfdrc6bPJxYZ5)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a})))$

Definition 4 We define $c_2Ebool_2E_2EF$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 7 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 8 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.(ap (c_2Emin_2E_40 (2^{A_27a}))$

Definition 9 We define $c_2Ecombin_2EUPDATE$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0a \in A_27a.\lambda V1b \in A_27b.(ap (c_2Emin_2E_40 (2^{A_27a}))$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \tag{2}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \tag{3}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{4}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\
& \quad \forall V0f \in (A_27b^{A_27a}). (\forall V1a \in A_27a. (\forall V2b \in A_27b. \\
& \quad ((ap\ V0f\ V1a) = V2b) \Leftrightarrow ((ap\ (ap\ (ap\ (c_2Ecombin_2EUPDATE\ A_27a\ A_27b) \\
& \quad \quad V1a)\ V2b)\ V0f) = V0f))))))
\end{aligned} \tag{5}$$

Theorem 1

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\
& \quad \forall V0f \in (A_27b^{A_27a}). (\forall V1a \in A_27a. ((ap\ (ap\ (ap\ (c_2Ecombin_2EUPDATE \\
& \quad \quad A_27a\ A_27b)\ V1a)\ (ap\ V0f\ V1a))\ V0f) = V0f)))
\end{aligned}$$