

thm_2Ecombin_2ESAME__KEY__UPDATE__DIFFER (TMWHD5X2RBVSyjvtHcaGvPBx3NYRyBr6ju8)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 8 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$.

Definition 9 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.(ap$

Definition 10 We define $c_2Ecombin_2EUPDATE$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0a \in A_27a.\lambda V1b \in A_27b$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \tag{1}$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \tag{2}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \tag{3}$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\ \forall V0f \in (A_27b^{A_27a}).(\forall V1g \in (A_27b^{A_27a}).((V0f = \\ V1g) \Leftrightarrow (\forall V2x \in A_27a.((ap\ V0f\ V2x) = (ap\ V1g\ V2x)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t1 \in A_27a.(\forall V1t2 \in \\ A_27a.(((ap\ (ap\ (ap\ (c_2Ebool_2ECOND\ A_27a)\ c_2Ebool_2ET)\ V0t1) \\ V1t2) = V0t1) \wedge ((ap\ (ap\ (ap\ (c_2Ebool_2ECOND\ A_27a)\ c_2Ebool_2EF) \\ V0t1)\ V1t2) = V1t2)))))) \end{aligned} \quad (5)$$

Theorem 1

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow \forall A_27c. \\ nonempty\ A_27c \Rightarrow \forall A_27d.nonempty\ A_27d \Rightarrow (\forall V0f \in (A_27d^{A_27c}). \\ (\forall V1f1 \in A_27a.(\forall V2f2 \in A_27b.(\forall V3a \in A_27c. \\ (\forall V4b \in A_27d.(\forall V5c \in A_27d.((\neg(V4b = V5c)) \Rightarrow (\neg((ap \\ (ap\ (ap\ (c_2Ecombin_2EUPDATE\ A_27c\ A_27d)\ V3a)\ V4b)\ V0f) = (ap\ (ap \\ (ap\ (c_2Ecombin_2EUPDATE\ A_27c\ A_27d)\ V3a)\ V5c)\ V0f)))))))))) \end{aligned}$$