

thm_2Ecombin_2EUPDATE__APPLY
(TMMuicaCzkRr3JkX2Qf4bzwjJNjj1bQ6vFw)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_EF$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_EF$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t$

Definition 8 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$.

Definition 9 We define $c_2Ebool_2E_COND$ to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.(ap$

Definition 10 We define $c_2Ecombin_2EUPDATE$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0a \in A_27a.\lambda V1b \in A_27b$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \tag{3}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (4)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t1 \in A_27a. (\forall V1t2 \in \\ & A_27a. (((ap\ (ap\ (ap\ (c_2Ebool_2ECOND\ A_27a)\ c_2Ebool_2ET)\ V0t1)\ V1t2) = V0t1) \wedge \\ & ((ap\ (ap\ (ap\ (c_2Ebool_2ECOND\ A_27a)\ c_2Ebool_2EF)\ V0t1)\ V1t2) = V1t2)))))) \end{aligned} \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow \forall A_27c. \\ & nonempty\ A_27c \Rightarrow \forall A_27d.nonempty\ A_27d \Rightarrow ((\forall V0a \in A_27a. \\ & (\forall V1x \in A_27b. (\forall V2f \in (A_27b^{A_27a}). ((ap\ (ap\ (ap\ (ap \\ & (c_2Ecombin_2EUPDATE\ A_27a\ A_27b)\ V0a)\ V1x)\ V2f)\ V0a) = V1x)))))) \wedge \\ & (\forall V3a \in A_27c. (\forall V4b \in A_27c. (\forall V5x \in A_27d. (\\ & \forall V6f \in (A_27d^{A_27c}). ((\neg(V3a = V4b)) \Rightarrow ((ap\ (ap\ (ap\ (ap\ (c_2Ecombin_2EUPDATE \\ & A_27c\ A_27d)\ V3a)\ V5x)\ V6f)\ V4b) = (ap\ V6f\ V4b)))))))))) \end{aligned}$$