

thm_2Ecombin_2EUPDATE__APPLY__ID
(TMTY83kwjMJv7dvfeQb9913eDVXBoAHZc93)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow p \ Q)$ of type ι .

Definition 2 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define `c_2Ebool_2E_2ET` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

Definition 4 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 5 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

Definition 6 We define `c_2Ebool_2E_2EF` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V0t \in 2. V0t)$.

Definition 7 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t)) (\text{c_2Ebool_2E_2EF } V0t)))$

Definition 8 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

Definition 9 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p \ (\text{ap } P \ x)) \ \text{then } (\text{the } (\lambda x. x \in A \wedge p \ (\text{ap } P \ x)))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define `c_2Ebool_2ECOND` to be $\lambda A. 27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A. 27a. (\lambda V2t2 \in A. 27a. (V2t2 \Rightarrow V1t1))))$

Definition 11 We define `c_2Ecombin_2EUPDATE` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0a \in A. 27a. \lambda V1b \in A. 27b. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0a)) (\text{c_2Emin_2E_3D_3D_3E } V1b))$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p \ V0t1) \Rightarrow (p \ V1t2)) \Rightarrow (((p \ V1t2) \Rightarrow (p \ V0t1)) \Rightarrow ((p \ V0t1) \Leftrightarrow (p \ V1t2)))))) \tag{2}$$

Assume the following.

$$(\forall V0t \in 2. (\text{False} \Rightarrow (p \ V0t))) \tag{3}$$

Assume the following.

$$(\forall V0t \in 2.((p V0t) \vee (\neg(p V0t)))) \quad (4)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (5)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0f \in (A_27b^{A_27a}).(\forall V1g \in (A_27b^{A_27a}).((V0f = V1g) \Leftrightarrow (\forall V2x \in A_27a.((ap\ V0f\ V2x) = (ap\ V1g\ V2x)))))) \quad (6)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (7)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t1 \in A_27a.(\forall V1t2 \in A_27a.(((ap\ (ap\ (ap\ (c_2Ebool_2ECOND\ A_27a)\ c_2Ebool_2ET)\ V0t1)\ V1t2) = V0t1) \wedge ((ap\ (ap\ (ap\ (c_2Ebool_2ECOND\ A_27a)\ c_2Ebool_2EF)\ V0t1)\ V1t2) = V1t2)))) \quad (8)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0f \in (A_27b^{A_27a}).(\forall V1a \in A_27a.(\forall V2b \in A_27b.(((ap\ V0f\ V1a) = V2b) \Leftrightarrow ((ap\ (ap\ (ap\ (c_2Ecombin_2EUPDATE\ A_27a\ A_27b)\ V1a)\ V2b)\ V0f) = V0f))))))$$