

thm_2Ecomplex_2ECOMPLETEX (TMVSVgC4Q9gyjHiTVYtXX4pScqGdJXNfxzv)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $c_2Epair_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EFST\ A_27a\ A_27b \in (A_27a^{(ty_2Epair_2Eprod\ A_27a\ A_27b)}) \tag{3}$$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ecomplex_2ERE$ to be $\lambda V0z \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)$

Let $c_2Epair_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2ESND\ A_27a\ A_27b \in (A_27b^{(ty_2Epair_2Eprod\ A_27a\ A_27b)}) \tag{4}$$

Definition 5 We define $c_2Ecomplex_2EIM$ to be $\lambda V0z \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)$

Definition 6 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21\ 2)) (\lambda V2t \in 2.V2t)))$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EABS_prod \\ A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ A_27b)^{(2^{A_27b})^{A_27a}}) \end{aligned} \quad (5)$$

Definition 8 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0x \in A_27a. \lambda V1y \in A_27b. (ap\ (c_2E$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in \\ A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow \\ True)) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\ \forall V0x \in (ty_2Epair_2Eprod\ A_27a\ A_27b). ((ap\ (ap\ (c_2Epair_2E_2C \\ A_27a\ A_27b)\ (ap\ (c_2Epair_2EFST\ A_27a\ A_27b)\ V0x))\ (ap\ (c_2Epair_2ESND \\ A_27a\ A_27b)\ V0x)) = V0x)) \end{aligned} \quad (9)$$

Theorem 1

$$\begin{aligned} (\forall V0z \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal). \\ ((ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal) \\ (ap\ c_2Ecomplex_2ERE\ V0z))\ (ap\ c_2Ecomplex_2EIM\ V0z)) = V0z)) \end{aligned}$$