

thm\_2Ecomplex\_2ECOMPLEX\_\_10  
(TMKJQZbzvA7GDNjhGv5fS8PTdAGh2CYRtRi)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})))$

**Definition 5** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

Let  $ty\_2Erealx\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealx\_2Ereal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $c\_2Epair\_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epair\_2ESND\ A\_27a\ A\_27b \in (A\_27b^{(ty\_2Epair\_2Eprod\ A\_27a\ A\_27b)}) \tag{3}$$

**Definition 6** We define  $c\_2Ecomplex\_2EIM$  to be  $\lambda V0z \in (ty\_2Epair\_2Eprod\ ty\_2Erealx\_2Ereal\ ty\_2Erealx\_2Ereal)$

Let  $c\_2Epair\_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epair\_2EFST\ A\_27a\ A\_27b \in (A\_27a^{(ty\_2Epair\_2Eprod\ A\_27a\ A\_27b)}) \tag{4}$$

**Definition 7** We define  $c\_2Ecomplex\_2ERE$  to be  $\lambda V0z \in (ty\_2Epair\_2Eprod\ ty\_2Erealx\_2Ereal\ ty\_2Erealx\_2Ereal)$

**Definition 8** We define  $c\_Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_Ebool\_2E\_21 2) (\lambda V2t \in 2)))$   
Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{5}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{6}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{7}$$

**Definition 9** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}) \tag{8}$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epair\_2EABS\_prod\ A\_27a\ A\_27b \in ((ty\_2Epair\_2Eprod\ A\_27a\ A\_27b)^{(2^{A\_27b})^{A\_27a}}) \tag{9}$$

**Definition 10** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2Epair\_2EABS\_prod\ A\_27a\ A\_27b)\ x\ y)$

**Definition 11** We define  $c\_2Ecomplex\_2Ecomplex\_of\_real$  to be  $\lambda V0x \in ty\_2Erealax\_2Ereal.(ap (ap (c\_2Epair\_2EABS\_prod\ x\ x)\ x\ x))$

**Definition 12** We define  $c\_2Ecomplex\_2Ecomplex\_of\_num$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap\ c\_2Ecomplex\_2Ecomplex\_of\_real\ n)$

**Definition 13** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{10}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{11}$$

**Definition 14** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num\ m)$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{12}$$

**Definition 15** We define  $c\_2Earithmetic\_2EBIT1$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap (ap\ c\_2Earithmetic\_2E\_2B\ n)\ n)$

**Definition 16** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

**Definition 17** We define  $c\_Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_7E$

Assume the following.

$$True \tag{13}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{14}$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \tag{15}$$

Assume the following.

$$(\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \tag{16}$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \tag{17}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \tag{18}$$

Assume the following.

$$(\forall V0z \in (ty\_2Epair\_2Eprod ty\_2Erealx\_2Ereal ty\_2Erealx\_2Ereal). (\forall V1w \in (ty\_2Epair\_2Eprod ty\_2Erealx\_2Ereal ty\_2Erealx\_2Ereal). ((V0z = V1w) \Leftrightarrow (((ap c\_2Ecomplex\_2ERE V0z) = (ap c\_2Ecomplex\_2ERE V1w)) \wedge ((ap c\_2Ecomplex\_2EIM V0z) = (ap c\_2Ecomplex\_2EIM V1w)))))) \tag{19}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27b.((ap (c\_2Epair\_2EFST A\_27a A\_27b) (ap (ap (c\_2Epair\_2E\_2C A\_27a A\_27b) V0x) V1y)) = V0x))) \tag{20}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27b.((ap (c\_2Epair\_2ESND A\_27a A\_27b) (ap (ap (c\_2Epair\_2E\_2C A\_27a A\_27b) V0x) V1y)) = V1y))) \tag{21}$$

Assume the following.

$$(\neg((ap c\_2Ereal\_2Ereal\_of\_num (ap c\_2Earithmetic\_2ENUMERAL (ap c\_2Earithmetic\_2EBIT1 c\_2Earithmetic\_2EZERO))) = (ap c\_2Ereal\_2Ereal\_of\_num c\_2Enum\_2E0))) \tag{22}$$

**Theorem 1**

$$\begin{aligned} & (\neg((ap\ c\_2Ecomplex\_2Ecomplex\_of\_num\ (ap\ c\_2Earithmetic\_2ENUMERAL \\ & (ap\ c\_2Earithmetic\_2EBIT1\ c\_2Earithmetic\_2EZERO)))) = (ap\ c\_2Ecomplex\_2Ecomplex\_of\_num \\ & \qquad\qquad\qquad c\_2Enum\_2E0))) \end{aligned}$$