

thm_2Ecomplex_2ECOMPLETE_EQ_RADD (TMbkhsjc1smr4jHviahXSqyB2FvK5uFC1pd)

October 26, 2020

Definition 1 We define c_2Emin_2E3D to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $c_2Epair_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow c_2Epair_2ESND\ A.27a\ A.27b \in (A.27b^{(ty_2Epair_2Eprod\ A.27a\ A.27b)}) \tag{3}$$

Definition 3 We define c_2Ebool_2E21 to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E3D (2^{A-27a}))))$

Definition 4 We define $c_2Ecomplex_2EIM$ to be $\lambda V0z \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{4}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal}) \tag{5}$$

Definition 5 We define c_2Emin_2E40 to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap\ P\ x)) then (the (\lambda x.x \in A \wedge p (ap\ P\ x)))$ of type $\iota \Rightarrow \iota$.

Definition 6 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E40 (ty_2Erealax_2Ereal_add : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_add \in (((ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)) (6)$$

Let $c_2Erealax_2Ereal_eq : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_eq \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)) (7)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)} (8)$$

Definition 7 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)$

Definition 8 We define $c_2Erealax_2Ereal_add$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Let $c_2Epair_2EFAST : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EFAST A_27a A_27b \in (A_27a)^{(ty_2Epair_2Eprod A_27a A_27b)} (9)$$

Definition 9 We define $c_2Ecomplex_2ERE$ to be $\lambda V0z \in (ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2Erealax_2Ereal)$

Definition 10 We define $c_2Emin_2E3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow p Q)$ of type ι .

Definition 11 We define $c_2Ebool_2E2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E21 2) (\lambda V2t \in 2)))$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EABS_prod A_27a A_27b \in ((ty_2Epair_2Eprod A_27a A_27b)^{(2^{A_27b} A_27a)}) (10)$$

Definition 12 We define c_2Epair_2E2C to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2Epair_2EABS_prod$

Definition 13 We define $c_2Ecomplex_2Ecomplex_add$ to be $\lambda V0z \in (ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2Erealax_2Ereal)$

Assume the following.

$$True (11)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) (12)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (13)$$

Assume the following.

$$\begin{aligned} & (\forall V0z \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)). \\ & (\forall V1w \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)). \\ & (((ap\ (ap\ c_2Ecomplex_2Ecomplex_add\ V0z)\ V1w) = (ap\ (ap\ c_2Ecomplex_2Ecomplex_add \\ & \quad V1w)\ V0z)))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} & (\forall V0z \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)). \\ & (\forall V1w \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)). \\ & (\forall V2v \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)). \\ & (((ap\ (ap\ c_2Ecomplex_2Ecomplex_add\ V0z)\ V1w) = (ap\ (ap\ c_2Ecomplex_2Ecomplex_add \\ & \quad V0z)\ V2v)) \Leftrightarrow (V1w = V2v)))) \end{aligned} \quad (15)$$

Theorem 1

$$\begin{aligned} & (\forall V0z \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)). \\ & (\forall V1w \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)). \\ & (\forall V2v \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)). \\ & (((ap\ (ap\ c_2Ecomplex_2Ecomplex_add\ V0z)\ V2v) = (ap\ (ap\ c_2Ecomplex_2Ecomplex_add \\ & \quad V1w)\ V2v)) \Leftrightarrow (V0z = V1w)))) \end{aligned}$$