

thm_2Ecomplex_2ECOMPLETE_RE_IM_EQ (TMa6BSzVycVMbgicdwz3X5s3bajiiQHHgoB)

October 26, 2020

Definition 1 We define c_2Emin_2E3D to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $c_2Epair_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EFST\ A_27a\ A_27b \in (A_27a\ A_27b)^{(ty_2Epair_2Eprod\ A_27a\ A_27b)} \tag{3}$$

Definition 3 We define c_2Ebool_2E21 to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E3D (2^{A_27a}))$

Definition 4 We define $c_2Ecomplex_2ERE$ to be $\lambda V0z \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)$

Let $c_2Epair_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2ESND\ A_27a\ A_27b \in (A_27b)^{(ty_2Epair_2Eprod\ A_27a\ A_27b)} \tag{4}$$

Definition 5 We define $c_2Ecomplex_2EIM$ to be $\lambda V0z \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)$

Definition 6 We define $c_2Emin_2E3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 7 We define $c_2Ebool_2E2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E21\ 2)) (\lambda V2t \in 2.V2t)))$

Assume the following.

$$True \quad (5)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A.27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (6)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0x \in A.27a.((V0x = V0x) \Leftrightarrow True)) \quad (7)$$

Assume the following.

$$\begin{aligned} &\forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow (\\ &\forall V0p \in (ty.2Epair.2Eprod\ A.27a\ A.27b).(\forall V1q \in (ty.2Epair.2Eprod \\ &A.27a\ A.27b).((V0p = V1q) \Leftrightarrow (((ap\ (c.2Epair.2EFST\ A.27a\ A.27b)\ V0p) = \\ &(ap\ (c.2Epair.2EFST\ A.27a\ A.27b)\ V1q)) \wedge ((ap\ (c.2Epair.2ESND\ A.27a \\ &A.27b)\ V0p) = (ap\ (c.2Epair.2ESND\ A.27a\ A.27b)\ V1q)))))) \quad (8) \end{aligned}$$

Theorem 1

$$\begin{aligned} &(\forall V0z \in (ty.2Epair.2Eprod\ ty.2Erealax.2Ereal\ ty.2Erealax.2Ereal). \\ &(\forall V1w \in (ty.2Epair.2Eprod\ ty.2Erealax.2Ereal\ ty.2Erealax.2Ereal). \\ &((V0z = V1w) \Leftrightarrow (((ap\ c.2Ecomplex.2ERE\ V0z) = (ap\ c.2Ecomplex.2ERE \\ &V1w)) \wedge ((ap\ c.2Ecomplex.2EIM\ V0z) = (ap\ c.2Ecomplex.2EIM\ V1w)))))) \end{aligned}$$