

thm_2Ecomplex_2ECOMPLEX__SCALAR__LMUL__ZERO
 (TMXSe-
 fRFF7Z9rXezr4YfhmM9QVAaJQuG4MW)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 3 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{4}$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealx_2Ereal^{ty_2Enum_2Enum}) \tag{5}$$

Definition 4 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow p\ Q)$ of type ι .

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})))$

Definition 6 We define $c_Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_Ebool_2E_21 2) (\lambda V2t \in 2.$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Epair_2Eprod A0 A1) \quad (6)$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EABS_prod A_27a A_27b \in ((ty_2Epair_2Eprod A_27a A_27b)^{(2^{A_27b})^{A_27a}}) \quad (7)$$

Definition 7 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2E$

Definition 8 We define $c_2Ecomplex_2Ecomplex_of_real$ to be $\lambda V0x \in ty_2Erealax_2Ereal.(ap (ap (c_2E$

Definition 9 We define $c_2Ecomplex_2Ecomplex_of_num$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap c_2Ecompl$

Let $c_2Epair_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2ESND A_27a A_27b \in (A_27b^{(ty_2Epair_2Eprod A_27a A_27b)}) \quad (8)$$

Definition 10 We define $c_2Ecomplex_2EIM$ to be $\lambda V0z \in (ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2E$

Let $c_2Epair_2EFST : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EFST A_27a A_27b \in (A_27a^{(ty_2Epair_2Eprod A_27a A_27b)}) \quad (9)$$

Definition 11 We define $c_2Ecomplex_2ERE$ to be $\lambda V0z \in (ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2E$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty ty_2Ehreal_2Ehreal \quad (10)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal_REP_CLASS}) \quad (11)$$

Definition 12 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x) \text{ of type } \iota \Rightarrow \iota).$

Definition 13 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 (t$

Let $c_2Erealax_2Etrealmul : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealmul \in (((ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)))(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal) \quad (12)$$

Let $c_2Erealax_2Etrealeq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealeq \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)) \quad (13)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)}}) \quad (14)$$

Definition 14 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)$

Definition 15 We define $c_2Erealax_2Ereal_mul$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 16 We define $c_2Ecomplex_2Ecomplex_scalar_lmul$ to be $\lambda V0k \in ty_2Erealax_2Ereal.\lambda V1z \in ty_2Erealax_2Ereal$

Assume the following.

$$True \quad (15)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (16)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (17)$$

Assume the following.

$$(\forall V0x \in ty_2Erealax_2Ereal.((ap\ (ap\ c_2Erealax_2Ereal_mul\ (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0))\ V0x) = (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0))) \quad (18)$$

Theorem 1

$$(\forall V0z \in (ty_2Epair_2Eprod\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal).((ap\ (ap\ c_2Ecomplex_2Ecomplex_scalar_lmul\ (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0))\ V0z) = (ap\ c_2Ecomplex_2Ecomplex_of_num\ c_2Enum_2E0))))$$