

# thm\_2Ecomplex\_2ECONJ\_\_NEG (TMLfGMSZm- Fer7gCsbiK8QxgnsCKEEEEeA6hC)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let  $ty\_2Erealax\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealax\_2Ereal \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $c\_2Epair\_2ESND : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epair\_2ESND\ A\_27a\ A\_27b \in (A\_27b^{(ty\_2Epair\_2Eprod\ A\_27a\ A\_27b)}) \tag{3}$$

**Definition 3** We define  $c\_2Ebool\_2E21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ecomplex\_2EIM$  to be  $\lambda V0z \in (ty\_2Epair\_2Eprod\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal)$

Let  $ty\_2Ehreal\_2Ehreal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehreal\_2Ehreal \tag{4}$$

Let  $c\_2Erealax\_2Ereal\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Erealax\_2Ereal\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Ehreal\_2Ehreal\ ty\_2Ehreal\_2Ehreal)})^{ty\_2Erealax\_2Ereal}) \tag{5}$$

**Definition 5** We define  $c\_2Emin\_2E40$  to be  $\lambda A.\lambda P \in 2^A.if\ (\exists x \in A.p\ (ap\ P\ x))\ then\ (the\ (\lambda x.x \in A \wedge p\ x))$  of type  $\iota \Rightarrow \iota$ .



Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (13)$$

Assume the following.

$$\begin{aligned} &\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ &\quad \forall V0x \in A\_27a. (\forall V1y \in A\_27b. ((ap\ (c\_2Epair\_2EFST\ A\_27a\ A\_27b)\ (ap\ (ap\ (c\_2Epair\_2E\_2C\ A\_27a\ A\_27b)\ V0x)\ V1y)) = V0x))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} &\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ &\quad \forall V0x \in A\_27a. (\forall V1y \in A\_27b. ((ap\ (c\_2Epair\_2ESND\ A\_27a\ A\_27b)\ (ap\ (ap\ (c\_2Epair\_2E\_2C\ A\_27a\ A\_27b)\ V0x)\ V1y)) = V1y))) \end{aligned} \quad (15)$$

**Theorem 1**

$$\begin{aligned} &(\forall V0z \in (ty\_2Epair\_2Eprod\ ty\_2Erealax\_2Ereal\ ty\_2Erealax\_2Ereal). \\ &\quad ((ap\ c\_2Ecomplex\_2Econj\ (ap\ c\_2Ecomplex\_2Ecomplex\_neg\ V0z)) = \\ &\quad (ap\ c\_2Ecomplex\_2Ecomplex\_neg\ (ap\ c\_2Ecomplex\_2Econj\ V0z)))) \end{aligned}$$