

# thm\_2Edivides\_2ENOT\_\_PRIME\_\_1 (TMPGiK- FcUKHaZM88HjqmZs6F5W8KVSBrh4Q)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Earithmetic\_2E\_2A : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2A \in ((ty\_2Enum\_2Enum)^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum} \tag{2}$$

**Definition 7** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$  then  $(the (\lambda x.x \in A \wedge p))$  of type  $\iota \Rightarrow \iota$ .

**Definition 8** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40 A\_27a P)))$

**Definition 9** We define  $c\_2Edivides\_2Edivides$  to be  $\lambda V0a \in ty\_2Enum\_2Enum.\lambda V1b \in ty\_2Enum\_2Enum$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in omega \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum)^{omega} \tag{4}$$

**Definition 10** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 11** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (5)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (6)$$

**Definition 12** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum)^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum} \quad (7)$$

**Definition 13** We define  $c\_2Earithmetic\_2EBIT1$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap\ (ap\ c\_2Earithmetic$

**Definition 14** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

**Definition 15** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E$

**Definition 16** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 17** We define  $c\_2Edivides\_2Eprime$  to be  $\lambda V0a \in ty\_2Enum\_2Enum.(ap\ (ap\ c\_2Ebool\_2E\_2F\_5C$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee \\ & (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & ((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge \\ & ((\neg False) \Leftrightarrow True))) \end{aligned} \quad (11)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (12)$$

Assume the following.

$$\forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow (\forall V0x \in A_{27a}. (\forall V1y \in A_{27a}. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (13)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (14)$$

Assume the following.

$$(\forall V0x \in 2. (\forall V1x_{27} \in 2. (\forall V2y \in 2. (\forall V3y_{27} \in 2. (((((p V0x) \Leftrightarrow (p V1x_{27})) \wedge ((p V1x_{27}) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_{27})))) \Rightarrow (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_{27}) \Rightarrow (p V3y_{27})))))))) \Rightarrow (15)$$

**Theorem 1**

$$(\neg (p (\text{ap } c_{2Edivides\_2Eprime} (\text{ap } c_{2Earithmetic\_2ENUMERAL} (\text{ap } c_{2Earithmetic\_2EBIT1} c_{2Earithmetic\_2EZERO}))))))$$