

thm_2Eextreal_2Einf__min
(TMVg6x6s5aN227xpSeVEiW4ARE7woDoDUQf)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Let $ty_2Eextreal_2Eextreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Eextreal_2Eextreal \tag{1}$$

Let $c_2Eextreal_2Eextreal_le : \iota$ be given. Assume the following.

$$c_2Eextreal_2Eextreal_le \in ((2^{ty_2Eextreal_2Eextreal})^{ty_2Eextreal_2Eextreal}) \tag{2}$$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $c_2Eextreal_2Eextreal_ainv : \iota$ be given. Assume the following.

$$c_2Eextreal_2Eextreal_ainv \in (ty_2Eextreal_2Eextreal)^{ty_2Eextreal_2Eextreal} \tag{3}$$

Definition 8 We define $c_2Ebool_2E_2IN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{4}$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EABS_prod\ A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ A_27b)^{(2^{A_27b})^{A_27a}}) \quad (5)$$

Definition 9 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap\ (c_2E$

Let $c_2Epred_set_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epred_set_2EGSPEC\ A_27a\ A_27b \in ((2^{A_27a})^{(ty_2Epair_2Eprod\ A_27a\ 2)^{A_27b}}) \quad (6)$$

Definition 10 We define $c_2Epred_set_2EIMAGE$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (A_27b^{A_27a}).\lambda V1s \in$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \quad (7)$$

Let $c_2Eextreal_2ENormal : \iota$ be given. Assume the following.

$$c_2Eextreal_2ENormal \in (ty_2Eextreal_2Eextreal^{ty_2Erealax_2Ereal}) \quad (8)$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (9)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal}) \quad (10)$$

Definition 11 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ if $(\exists x \in A.p\ (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 12 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E_40\ (t$

Let $c_2Erealax_2Etreallt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreallt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (11)$$

Definition 13 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.$

Definition 14 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ V0P\ (ap\ (c_2Emin_2E_40\ (t$

Definition 15 We define c_2Ereal_2Esup to be $\lambda V0P \in (2^{ty_2Erealax_2Ereal}).(ap\ (c_2Emin_2E_40\ ty_2Ereal$

Let $c_2Eextreal_2ENegInf : \iota$ be given. Assume the following.

$$c_2Eextreal_2ENegInf \in ty_2Eextreal_2Eextreal \quad (12)$$

Assume the following.

$$\begin{aligned}
& (\forall V0p \in (2^{ty_2Eextreal_2Eextreal}).(\forall V1x \in ty_2Eextreal_2Eextreal. \\
& (((ap\ c_2Eextreal_2Eextreal_inf\ V0p) = V1x) \Leftrightarrow ((\forall V2y \in ty_2Eextreal_2Eextreal. \\
& ((p\ (ap\ V0p\ V2y)) \Rightarrow (p\ (ap\ (ap\ c_2Eextreal_2Eextreal_le\ V1x)\ V2y)))) \wedge \\
& (\forall V3y \in ty_2Eextreal_2Eextreal.((\forall V4z \in ty_2Eextreal_2Eextreal. \\
& ((p\ (ap\ V0p\ V4z)) \Rightarrow (p\ (ap\ (ap\ c_2Eextreal_2Eextreal_le\ V3y)\ V4z)))) \Rightarrow \\
& (p\ (ap\ (ap\ c_2Eextreal_2Eextreal_le\ V3y)\ V1x))))))
\end{aligned} \tag{22}$$

Theorem 1

$$\begin{aligned}
& (\forall V0p \in (2^{ty_2Eextreal_2Eextreal}).(\forall V1z \in ty_2Eextreal_2Eextreal. \\
& (((p\ (ap\ V0p\ V1z)) \wedge (\forall V2x \in ty_2Eextreal_2Eextreal.((p\ (\\
& ap\ V0p\ V2x)) \Rightarrow (p\ (ap\ (ap\ c_2Eextreal_2Eextreal_le\ V1z)\ V2x)))))) \Rightarrow \\
& ((ap\ c_2Eextreal_2Eextreal_inf\ V0p) = V1z)))
\end{aligned}$$