

thm_2Eextreal_2Emax_refl (TMN- HdASDu4BeFed1uXm9AfVJPVn36wnxtfH)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2ET` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2))) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x)$

Let `ty_2Eextreal_2Eextreal` : ι be given. Assume the following.

$$\text{nonempty } \text{ty_2Eextreal_2Eextreal} \tag{1}$$

Let `c_2Eextreal_2Eextreal_le` : ι be given. Assume the following.

$$\text{c_2Eextreal_2Eextreal_le} \in ((2^{\text{ty_2Eextreal_2Eextreal}})^{\text{ty_2Eextreal_2Eextreal}}) \tag{2}$$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. \lambda 27a : \iota. (\lambda V0P \in (2^{A \cdot 27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A \cdot 27a}))))$

Definition 4 We define `c_2Ebool_2EF` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V0t \in 2. V0t)$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

Definition 7 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p (\text{ap } P x)))$ of type $\iota \Rightarrow \iota$.

Definition 8 We define `c_2Ebool_2ECOND` to be $\lambda A. \lambda 27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A. 27a. (\lambda V2t2 \in A. 27a. (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } (V1t1)) (V2t2))))$

Definition 9 We define `c_2Eextreal_2Eextreal_max` to be $\lambda V0x \in \text{ty_2Eextreal_2Eextreal}. \lambda V1y \in \text{ty_2Eextreal_2Eextreal}. \text{max } (V0x) (V1y)$

Assume the following.

$$\text{True} \tag{3}$$

Assume the following.

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow (\forall V0x \in A. 27a. ((V0x = V0x) \Leftrightarrow \text{True})) \tag{4}$$

Assume the following.

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow (\forall V0b \in 2. (\forall V1t \in A. 27a. ((\text{ap } (\text{ap } (\text{ap } (\text{c_2Ebool_2ECOND } A. 27a) V0b) V1t) V1t) = V1t))) \tag{5}$$

Theorem 1

$$(\forall V0x \in ty_2Eextreal_2Eextreal.((ap (ap c_2Eextreal_2Eextreal_max V0x) V0x) = V0x))$$