

thm_2Efinite_map_2EDRESTRICT_FUNION_SAME (TMJKKpinjEjtJMj7gEHsjDLDognb5EZWQte)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $ty_2Efinite_map_2E fmap : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Efinite_map_2E fmap A0 A1) \quad (1)$$

Let $c_2Efinite_map_2EDRESTRICT : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Efinite_map_2EDRESTRICT A_27a A_27b \in (((ty_2Efinite_map_2E fmap A_27a A_27b)^{(2^{A_27a})})(ty_2Efinite_map_2E fmap A_27a A_27b)) \quad (2)$$

Let $c_2Efinite_map_2EFUNION : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Efinite_map_2EFUNION A_27a A_27b \in (((ty_2Efinite_map_2E fmap A_27a A_27b)^{(ty_2Efinite_map_2E fmap A_27a A_27b)}}(ty_2Efinite_map_2E fmap A_27a A_27b)) \quad (3)$$

Let $ty_2Eone_2Eone : \iota$ be given. Assume the following.

$$nonempty\ ty_2Eone_2Eone \quad (4)$$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Esum_2Esum\ A0\ A1) \quad (5)$$

Let $c_2Efinite_map_2Efmap_REP : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Efinite_map_2Efmap_REP\ A_27a\ A_27b \in ((ty_2Esum_2Esum\ A_27b\ ty_2Eone_2Eone)^{A_27a})^{(ty_2Efinite_map_2Efmap\ A_27a\ A_27b)} \quad (6)$$

Let $c_2Esum_2EOUTL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Esum_2EOUTL\ A_27a\ A_27b \in (A_27a)^{(ty_2Esum_2Esum\ A_27a\ A_27b)} \quad (7)$$

Definition 8 We define $c_2Efinite_map_2EFAPPLY$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (ty_2Efinite_map_2E$

Let $c_2Esum_2EISL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Esum_2EISL\ A_27a\ A_27b \in (2^{(ty_2Esum_2Esum\ A_27a\ A_27b)}) \quad (8)$$

Definition 9 We define $c_2Efinite_map_2EFDOM$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (ty_2Efinite_map_2E$

Definition 10 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(\lambda p\ V1f\ V0x)))$

Definition 11 We define $c_2Efinite_map_2ESUBMAP$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (ty_2Efinite_map_2E$

Assume the following.

$$True \quad (9)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (10)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (11)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0f \in (ty_2Efinite_map_2Efmap\ A_27a\ A_27b).(\forall V1P \in (2^{A_27a}).(p\ (ap\ (ap\ (c_2Efinite_map_2ESUBMAP\ A_27a\ A_27b)\ (ap\ (ap\ (c_2Efinite_map_2EDRESTRICT\ A_27a\ A_27b)\ V0f)\ V1P))\ V0f)))) \quad (12)$$

Assume the following.

$$\begin{aligned}
& \forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow \forall A_{.27b}.nonempty\ A_{.27b} \Rightarrow (\\
& \quad \forall V0f \in (ty_2Efinite_map_2Efm\ A_{.27a}\ A_{.27b}). (\forall V1g \in \\
& (ty_2Efinite_map_2Efm\ A_{.27a}\ A_{.27b}). ((p\ (ap\ (ap\ (c_2Efinite_map_2ESUBMAP \\
& \quad A_{.27a}\ A_{.27b})\ V0f)\ V1g)) \Leftrightarrow ((ap\ (ap\ (c_2Efinite_map_2EFUNION\ A_{.27a} \\
& \quad A_{.27b})\ V0f)\ V1g) = V1g)))) \\
& \hspace{15em} (13)
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& \forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow \forall A_{.27b}.nonempty\ A_{.27b} \Rightarrow (\\
& \quad \forall V0fm \in (ty_2Efinite_map_2Efm\ A_{.27a}\ A_{.27b}). (\forall V1s \in \\
& (2^{A_{.27a}}). ((ap\ (ap\ (c_2Efinite_map_2EFUNION\ A_{.27a}\ A_{.27b})\ (ap \\
& (ap\ (c_2Efinite_map_2EDRESTRICT\ A_{.27a}\ A_{.27b})\ V0fm)\ V1s))\ V0fm) = \\
& \quad V0fm)))
\end{aligned}$$