

thm_2Efinite_map_2EFDOM_EMPTY (TMD- wqsjtyJzofyDWdWLR8EtmWGzU5TvCoAu)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2ET` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. \lambda 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 4 We define `c_2Ebool_2E_2EF` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow P \Rightarrow Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V2t \in 2.V2t))))$

Definition 7 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 8 We define `c_2Ebool_2ECOND` to be $\lambda A. \lambda 27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A. 27a. (\lambda V2t2 \in A. 27a. (\text{ap } (\text{c_2Emin_2E_40 } A) V2t2))))$

Definition 9 We define `c_2Ebool_2E_3F` to be $\lambda A. \lambda 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } V0P (\text{ap } (\text{c_2Emin_2E_40 } A) V0P))))$

Definition 10 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V2t \in 2.V2t))))$

Let `ty_2Eone_2Eone` : ι be given. Assume the following.

$$\text{nonempty } \text{ty_2Eone_2Eone} \tag{1}$$

Let `ty_2Esum_2Esum` : $\iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. \text{nonempty } A0 \Rightarrow \forall A1. \text{nonempty } A1 \Rightarrow \text{nonempty } (\text{ty_2Esum_2Esum } A0 A1) \tag{2}$$

Let `c_2Esum_2EABS_sum` : $\iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow \forall A. 27b. \text{nonempty } A. 27b \Rightarrow \text{c_2Esum_2EABS_sum } A. 27a A. 27b \in ((\text{ty_2Esum_2Esum } A. 27a A. 27b)^{((2^{A-27b})^{A-27a})^2}) \tag{3}$$

Definition 11 We define c_Esum_2EINL to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0e \in A_27a. (ap (c_Esum_2EABS$

Definition 12 We define c_Eone_2Eone to be $(ap (c_Emin_2E_40 ty_2Eone_2Eone) (\lambda V0x \in ty_2Eone_2E$

Definition 13 We define $c_Ebool_2E_7E$ to be $(\lambda V0t \in 2. (ap (ap c_Emin_2E_3D_3D_3E V0t) c_Ebool_2E$

Definition 14 We define c_Esum_2EINR to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0e \in A_27b. (ap (c_Esum_2EABS$

Definition 15 We define $c_Efinite_map_2Eis_fmap$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. (\lambda V0a0 \in ((ty_2Esum_2E$

Let $ty_2Efinite_map_2Efmap : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. nonempty A0 \Rightarrow \forall A1. nonempty A1 \Rightarrow nonempty (ty_2Efinite_map_2Efmap A0 A1) \quad (4)$$

Let $c_2Efinite_map_2Efmap_ABS : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow c_2Efinite_map_2Efmap_ABS A_27a A_27b \in ((ty_2Efinite_map_2Efmap A_27a A_27b)^{(ty_2Esum_2Esum A_27b ty_2Eone_2Eone)^{A_27a}}) \quad (5)$$

Definition 16 We define $c_2Efinite_map_2EFEMPTY$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. (ap (c_2Efinite_map_2E$

Let $c_2Efinite_map_2Efmap_REP : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow c_2Efinite_map_2Efmap_REP A_27a A_27b \in (((ty_2Esum_2Esum A_27b ty_2Eone_2Eone)^{A_27a})^{(ty_2Efinite_map_2Efmap A_27a A_27b)}) \quad (6)$$

Let $c_2Esum_2EISL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow c_2Esum_2EISL A_27a A_27b \in (2^{(ty_2Esum_2Esum A_27a A_27b)}) \quad (7)$$

Definition 17 We define $c_2Efinite_map_2EFDOM$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0f \in (ty_2Efinite_map$

Definition 18 We define $c_2Epred_set_2EEMPTY$ to be $\lambda A_27a : \iota. (\lambda V0x \in A_27a. c_2Ebool_2EF)$.

Definition 19 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota. (\lambda V0x \in A_27a. (\lambda V1f \in (2^{A_27a}). (ap V1f V0x)))$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p V0t))) \quad (10)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0t1 \in A_27a. (\forall V1t2 \in A_27b. ((\lambda V2x \in A_27b. V0t1\ V1t2) = V0t1))) \quad (11)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (12)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (13)$$

Assume the following.

$$((\forall V0t \in 2. ((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (14)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (15)$$

Assume the following.

$$(\forall V0x \in 2. (\forall V1y \in 2. (\forall V2z \in 2. (\forall V3w \in 2. (((p\ V0x) \Rightarrow (p\ V1y)) \wedge ((p\ V2z) \Rightarrow (p\ V3w))) \Rightarrow (((p\ V0x) \wedge (p\ V2z)) \Rightarrow ((p\ V1y) \wedge (p\ V3w)))))) \quad (16)$$

Assume the following.

$$(\forall V0x \in 2. (\forall V1y \in 2. (\forall V2z \in 2. (\forall V3w \in 2. (((p\ V0x) \Rightarrow (p\ V1y)) \wedge ((p\ V2z) \Rightarrow (p\ V3w))) \Rightarrow (((p\ V0x) \vee (p\ V2z)) \Rightarrow ((p\ V1y) \vee (p\ V3w)))))) \quad (17)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1Q \in (2^{A_27a}). ((\forall V2x \in A_27a. ((p\ (ap\ V0P\ V2x)) \Rightarrow (p\ (ap\ V1Q\ V2x)))) \Rightarrow ((\exists V3x \in A_27a. (p\ (ap\ V0P\ V3x))) \Rightarrow (\exists V4x \in A_27a. (p\ (ap\ V1Q\ V4x)))))) \quad (18)$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\
& (\forall V0a \in (ty_2Efinite_map_2E fmap\ A_27a\ A_27b).((ap\ (c_2Efinite_map_2E fmap_ABS \\
& \quad A_27a\ A_27b)\ (ap\ (c_2Efinite_map_2E fmap_REP\ A_27a\ A_27b)\ V0a)) = \\
& \quad V0a)) \wedge (\forall V1r \in ((ty_2Esum_2Esum\ A_27b\ ty_2Eone_2Eone)^{A_27a}). \\
& \quad ((p\ (ap\ (c_2Efinite_map_2Eis_fmap\ A_27a\ A_27b)\ V1r)) \Leftrightarrow ((ap\ (\\
& \quad c_2Efinite_map_2E fmap_REP\ A_27a\ A_27b)\ (ap\ (c_2Efinite_map_2E fmap_ABS \\
& \quad \quad A_27a\ A_27b)\ V1r))) = V1r)))) \\
& \hspace{15em} (19)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}).(\forall V1x \in \\
& A_27a.((p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27a)\ V1x)\ V0P)) \Leftrightarrow (p\ (ap\ V0P\ V1x)))))) \\
& \hspace{15em} (20)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0s \in (2^{A_27a}).(\forall V1t \in \\
& (2^{A_27a}).((V0s = V1t) \Leftrightarrow (\forall V2x \in A_27a.((p\ (ap\ (ap\ (c_2Ebool_2EIN \\
& \quad A_27a)\ V2x)\ V0s)) \Leftrightarrow (p\ (ap\ (ap\ (c_2Ebool_2EIN\ A_27a)\ V2x)\ V1t)))))) \\
& \hspace{15em} (21)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.(\neg(p\ (ap\ (ap \\
& \quad (c_2Ebool_2EIN\ A_27a)\ V0x)\ (c_2Epred_set_2EEMPTY\ A_27a)))))) \\
& \hspace{15em} (22)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\
& (\forall V0x \in A_27a.(p\ (ap\ (c_2Esum_2EISL\ A_27a\ A_27b)\ (ap\ (c_2Esum_2EINL \\
& \quad A_27a\ A_27b)\ V0x)))) \wedge (\forall V1y \in A_27b.(\neg(p\ (ap\ (c_2Esum_2EISL \\
& \quad A_27a\ A_27b)\ (ap\ (c_2Esum_2EINR\ A_27a\ A_27b)\ V1y)))))) \\
& \hspace{15em} (23)
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\
& (ap\ (c_2Efinite_map_2EFDOM\ A_27a\ A_27b)\ (c_2Efinite_map_2EEMPTY \\
& \quad A_27a\ A_27b)) = (c_2Epred_set_2EEMPTY\ A_27a))
\end{aligned}$$