

thm_2Efinite__map_2EFDOM__o__f (TM- bUSCdYoF3T4WmUAiSXLuP3G2VeSjUjo6R)

October 26, 2020

Let $ty_2Efinite_map_2Efdom_o_f : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Efinite_map_2Efdom_o_f\ A0\ A1) \quad (1)$$

Let $c_2Efinite_map_2Eo_f : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow \forall A.27c. \\ & nonempty\ A.27c \Rightarrow c_2Efinite_map_2Eo_f\ A.27a\ A.27b\ A.27c \in (((\\ & ty_2Efinite_map_2Efdom_o_f\ A.27a\ A.27c)^{(ty_2Efinite_map_2Efdom_o_f\ A.27a\ A.27b)})^{(A.27c^{A.27b})}) \end{aligned} \quad (2)$$

Let $ty_2Eone_2Eone : \iota$ be given. Assume the following.

$$nonempty\ ty_2Eone_2Eone \quad (3)$$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Esum_2Esum\ A0\ A1) \quad (4)$$

Let $c_2Efinite_map_2Efdom_REP : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow c_2Efinite_map_2Efdom_REP\ A.27a\ A.27b \in (((ty_2Esum_2Esum\ A.27b\ ty_2Eone_2Eone)^{A.27a})^{(ty_2Efinite_map_2Efdom_o_f\ A.27a\ A.27b)}) \quad (5)$$

Let $c_2Esum_2EISL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow c_2Esum_2EISL\ A.27a\ A.27b \in (2^{(ty_2Esum_2Esum\ A.27a\ A.27b)}) \quad (6)$$

Definition 1 We define c_2Emin_2E3D to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o\ (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_Ebool_2E_21$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap (ap (c_Emin_2E_3D (2^{A_27a})))$

Definition 4 We define $c_Efinite_map_2EFDOM$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0f \in (ty_2Efinite_map_2E$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow \forall A_27c. \\ & nonempty A_27c \Rightarrow (\forall V0f \in (A_27c^{A_27b}). (\forall V1g \in (ty_2Efinite_map_2E fmap \\ & A_27a A_27b). ((ap (c_Efinite_map_2EFDOM A_27a A_27b) V1g) = \\ & (ap (c_Efinite_map_2EFDOM A_27a A_27c) (ap (ap (c_2Efinite_map_2Eo_f \\ & A_27a A_27b A_27c) V0f) V1g)))))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow \forall A_27c. \\ & nonempty A_27c \Rightarrow (\forall V0f \in (A_27c^{A_27b}). (\forall V1g \in (ty_2Efinite_map_2E fmap \\ & A_27a A_27b). ((ap (c_Efinite_map_2EFDOM A_27a A_27c) (ap (ap \\ & (c_2Efinite_map_2Eo_f A_27a A_27b A_27c) V0f) V1g)) = (ap (c_Efinite_map_2EFDOM \\ & A_27a A_27b) V1g)))))) \end{aligned}$$