

thm\_2Efinite\_\_map\_2EFUNION\_\_IDEMPOT  
(TM-  
SaW3FvQj5FFLxCP7uPfbjeENkKoDd2FUj)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o(x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2)) (\lambda V0t \in 2.V0t)$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2. \lambda Q \in 2. inj\_o(p \Rightarrow Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2. (ap (ap (c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

**Definition 7** We define  $c\_2Ebool\_2E\_2IN$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. (\lambda V1f \in (2^{A\_27a}). (ap V1f V0x)))$

**Definition 8** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2)) (\lambda V2t \in 2.V2t)))$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A. \lambda P \in 2^A. \mathbf{if} (\exists x \in A. p (ap P x)) \mathbf{then} (the (\lambda x. x \in A \wedge p x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_2COND$  to be  $\lambda A\_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A\_27a. (\lambda V2t2 \in A\_27a. (ap V1t1 V2t2)))$

Let  $ty\_2Eone\_2Eone : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Eone\_2Eone \tag{1}$$

Let  $ty\_2Esum\_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0. nonempty\ A0 \Rightarrow \forall A1. nonempty\ A1 \Rightarrow nonempty\ (ty\_2Esum\_2Esum\ A0\ A1) \tag{2}$$

Let  $ty\_2Efinite\_map\_2E fmap : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Efinite\_map\_2E\ fmap\ A0\ A1)$$
(3)

Let  $c\_2Efinite\_map\_2E fmap\_REP : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Efinite\_map\_2E\ fmap\_REP\ A\_27a\ A\_27b \in (((ty\_2Esum\_2Esum\ A\_27b\ ty\_2Eone\_2Eone)^{A\_27a})^{(ty\_2Efinite\_map\_2E\ fmap\ A\_27a\ A\_27b)})$$
(4)

Let  $c\_2Esum\_2EOUTL : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EOUTL\ A\_27a\ A\_27b \in (A\_27a)^{(ty\_2Esum\_2Esum\ A\_27a\ A\_27b)}$$
(5)

**Definition 11** We define  $c\_2Efinite\_map\_2EFAPPLY$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0f \in (ty\_2Efinite\_map\_2EFUNION$

Let  $c\_2Efinite\_map\_2EFUNION : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Efinite\_map\_2EFUNION\ A\_27a\ A\_27b \in (((ty\_2Efinite\_map\_2E\ fmap\ A\_27a\ A\_27b)^{(ty\_2Efinite\_map\_2E\ fmap\ A\_27a\ A\_27b)})^{(ty\_2Efinite\_map\_2E\ fmap\ A\_27a\ A\_27b)})$$
(6)

Let  $c\_2Esum\_2EISL : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EISL\ A\_27a\ A\_27b \in (2^{(ty\_2Esum\_2Esum\ A\_27a\ A\_27b)})$$
(7)

**Definition 12** We define  $c\_2Efinite\_map\_2EFDOM$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0f \in (ty\_2Efinite\_map\_2EFUNION$

**Definition 13** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2E\ prod\ A0\ A1)$$
(8)

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epair\_2EABS\_prod\ A\_27a\ A\_27b \in ((ty\_2Epair\_2E\ prod\ A\_27a\ A\_27b)^{(2^{A\_27b})^{A\_27a}})$$
(9)

**Definition 14** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap\ (c\_2E$

Let  $c\_2Epred\_set\_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epred\_set\_2EGSPEC\ A\_27a\ A\_27b \in ((2^{A\_27a})^{(ty\_2Epair\_2E\ prod\ A\_27a\ 2)^{A\_27b}})$$
(10)

**Definition 15** We define  $c\_2Epred\_set\_2EUNION$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A\_27a}).\lambda V1t \in (2^{A\_27a}).(ap (c$

Assume the following.

$$True \quad (11)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (12)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \quad (13) \end{aligned}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (( \\ & (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (14) \end{aligned}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (15)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (16)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0b \in 2.(\forall V1t \in A\_27a.((ap (ap (ap (c\_2Ebool\_2ECOND\ A\_27a)\ V0b)\ V1t)\ V1t) = V1t))) \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p\ V0t1) \Rightarrow \\ & ((p\ V1t2) \Rightarrow (p\ V2t3))) \Leftrightarrow (((p\ V0t1) \wedge (p\ V1t2)) \Rightarrow (p\ V2t3)))))) \quad (18) \end{aligned}$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2.(\forall V1x\_27 \in 2.(\forall V2y \in 2.(\forall V3y\_27 \in \\ & 2.(((p\ V0x) \Leftrightarrow (p\ V1x\_27)) \wedge ((p\ V1x\_27) \Rightarrow ((p\ V2y) \Leftrightarrow (p\ V3y\_27)))) \Rightarrow \\ & (((p\ V0x) \Rightarrow (p\ V2y)) \Leftrightarrow ((p\ V1x\_27) \Rightarrow (p\ V3y\_27)))))) \quad (19) \end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow \forall A_{.27b}.nonempty\ A_{.27b} \Rightarrow ( \\
& \quad \forall V0f \in (ty\_2Efinite\_map\_2E fmap\ A_{.27a}\ A_{.27b}).(\forall V1g \in \\
& \quad (ty\_2Efinite\_map\_2E fmap\ A_{.27a}\ A_{.27b}).(((ap\ (c\_2Efinite\_map\_2EFDOM \\
& \quad A_{.27a}\ A_{.27b})\ V0f) = (ap\ (c\_2Efinite\_map\_2EFDOM\ A_{.27a}\ A_{.27b})\ V1g)) \wedge \\
& \quad (\forall V2x \in A_{.27a}.((p\ (ap\ (ap\ (c\_2Ebool\_2EIN\ A_{.27a})\ V2x)\ (ap\ ( \\
& \quad c\_2Efinite\_map\_2EFDOM\ A_{.27a}\ A_{.27b})\ V0f)))) \Rightarrow ((ap\ (ap\ (c\_2Efinite\_map\_2EFAPPLY \\
& \quad A_{.27a}\ A_{.27b})\ V0f)\ V2x) = (ap\ (ap\ (c\_2Efinite\_map\_2EFAPPLY\ A_{.27a} \\
& \quad A_{.27b})\ V1g)\ V2x)))))) \Leftrightarrow (V0f = V1g))
\end{aligned} \tag{20}$$

Assume the following.

$$\begin{aligned}
& \forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow \forall A_{.27b}.nonempty\ A_{.27b} \Rightarrow ( \\
& \quad \forall V0f \in (ty\_2Efinite\_map\_2E fmap\ A_{.27a}\ A_{.27b}).(\forall V1g \in \\
& \quad (ty\_2Efinite\_map\_2E fmap\ A_{.27a}\ A_{.27b}).((ap\ (c\_2Efinite\_map\_2EFDOM \\
& \quad A_{.27a}\ A_{.27b})\ (ap\ (ap\ (c\_2Efinite\_map\_2EFUNION\ A_{.27a}\ A_{.27b})\ V0f) \\
& \quad V1g)) = (ap\ (ap\ (c\_2Epred\_set\_2EUNION\ A_{.27a})\ (ap\ (c\_2Efinite\_map\_2EFDOM \\
& \quad A_{.27a}\ A_{.27b})\ V0f))\ (ap\ (c\_2Efinite\_map\_2EFDOM\ A_{.27a}\ A_{.27b})\ V1g))) \wedge \\
& \quad (\forall V2x \in A_{.27a}.((ap\ (ap\ (c\_2Efinite\_map\_2EFAPPLY\ A_{.27a} \\
& \quad A_{.27b})\ (ap\ (ap\ (c\_2Efinite\_map\_2EFUNION\ A_{.27a}\ A_{.27b})\ V0f)\ V1g)) \\
& \quad V2x) = (ap\ (ap\ (ap\ (c\_2Ebool\_2ECOND\ A_{.27b})\ (ap\ (ap\ (c\_2Ebool\_2EIN \\
& \quad A_{.27a})\ V2x)\ (ap\ (c\_2Efinite\_map\_2EFDOM\ A_{.27a}\ A_{.27b})\ V0f)))\ (ap \\
& \quad (ap\ (c\_2Efinite\_map\_2EFAPPLY\ A_{.27a}\ A_{.27b})\ V0f)\ V2x))\ (ap\ (ap\ ( \\
& \quad c\_2Efinite\_map\_2EFAPPLY\ A_{.27a}\ A_{.27b})\ V1g)\ V2x))))))
\end{aligned} \tag{21}$$

Assume the following.

$$\forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow (\forall V0s \in (2^{A_{.27a}}).((ap\ (ap\ (c\_2Epred\_set\_2EUNION\ A_{.27a})\ V0s)\ V0s) = V0s)) \tag{22}$$

**Theorem 1**

$$\begin{aligned}
& \forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow \forall A_{.27b}.nonempty\ A_{.27b} \Rightarrow ( \\
& \quad \forall V0fm \in (ty\_2Efinite\_map\_2E fmap\ A_{.27a}\ A_{.27b}).((ap\ (ap \\
& \quad (c\_2Efinite\_map\_2EFUNION\ A_{.27a}\ A_{.27b})\ V0fm)\ V0fm) = V0fm))
\end{aligned}$$