

thm_2Efinite__map_2ESUBMAP__FEMPTY
 (TMUwd-
 FuWf6JJ2oFJ4sYzXUaMaU4RbBNmCCC)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a})))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Let $ty_2Eone_2Eone : \iota$ be given. Assume the following.

$$nonempty\ ty_2Eone_2Eone \tag{1}$$

Definition 5 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p))$ of type $\iota \Rightarrow \iota$.

Definition 6 We define c_2Eone_2Eone to be $(ap (c_2Emin_2E_40 ty_2Eone_2Eone) (\lambda V0x \in ty_2Eone_2Eone))$

Definition 7 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 8 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Esum_2Esum\ A0\ A1) \tag{2}$$

Let $c_2Esum_2EABS_sum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Esum_2EABS_sum\ A_27a\ A_27b \in ((ty_2Esum_2Esum\ A_27a\ A_27b)^{((2^{A_27b})^{A_27a})^2}) \tag{3}$$

Definition 10 We define c_2Esum_2EINR to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0e \in A_27b. (ap (c_2Esum_2EABS$

Let $ty_2Efinite_map_2Efmap : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Efinite_map_2Efmap A0 A1) \quad (4)$$

Let $c_2Efinite_map_2Efmap_ABS : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Efinite_map_2Efmap_ABS A_27a A_27b \in ((ty_2Efinite_map_2Efmap A_27a A_27b)^{(ty_2Esum_2Esum A_27b ty_2Eone_2Eone)^{A_27a}}) \quad (5)$$

Definition 11 We define $c_2Efinite_map_2EFEMPTY$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. (ap (c_2Efinite_map_2E$

Let $c_2Efinite_map_2Efmap_REP : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Efinite_map_2Efmap_REP A_27a A_27b \in (((ty_2Esum_2Esum A_27b ty_2Eone_2Eone)^{A_27a})^{(ty_2Efinite_map_2Efmap A_27a A_27b)}) \quad (6)$$

Let $c_2Esum_2EOUTL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Esum_2EOUTL A_27a A_27b \in (A_27a^{(ty_2Esum_2Esum A_27a A_27b)}) \quad (7)$$

Definition 12 We define $c_2Efinite_map_2EFAPPLY$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0f \in (ty_2Efinite_map$

Let $c_2Esum_2EISL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Esum_2EISL A_27a A_27b \in (2^{(ty_2Esum_2Esum A_27a A_27b)}) \quad (8)$$

Definition 13 We define $c_2Efinite_map_2EFDOM$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0f \in (ty_2Efinite_map$

Definition 14 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota. (\lambda V0x \in A_27a. (\lambda V1f \in (2^{A_27a}). (ap V1f V0x)))$

Definition 15 We define $c_2Efinite_map_2ESUBMAP$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0f \in (ty_2Efinite_map$

Definition 16 We define $c_2Epred_set_2EEMPTY$ to be $\lambda A_27a : \iota. (\lambda V0x \in A_27a. c_2Ebool_2EF)$.

Assume the following.

$$True \quad (9)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (10)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p V0t))) \quad (11)$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a. (p V0t)) \Leftrightarrow (p V0t))) \quad (12)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \quad (13)$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (14)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p V0t1) \Rightarrow \\ & ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2. (\forall V1x_27 \in 2. (\forall V2y \in 2. (\forall V3y_27 \in \\ & 2. (((p V0x) \Leftrightarrow (p V1x_27)) \wedge ((p V1x_27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_27)))) \Rightarrow \\ & (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_27) \Rightarrow (p V3y_27)))))) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow (\\ & (ap (c_2Efinite_map_2EFDOM A_27a A_27b) (c_2Efinite_map_2EFEMPTY \\ & A_27a A_27b)) = (c_2Epred_set_2EEMPTY A_27a)) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & \forall A_27a. nonempty A_27a \Rightarrow (\forall V0x \in A_27a. (\neg (ap (ap \\ & (c_2Ebool_2EIN A_27a) V0x) (c_2Epred_set_2EEMPTY A_27a)))) \end{aligned} \quad (18)$$

Theorem 1

$$\begin{aligned} & \forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow (\\ & \forall V0f \in (ty_2Efinite_map_2E fmap A_27a A_27b). (p (ap (ap \\ & (c_2Efinite_map_2ESUBMAP A_27a A_27b) (c_2Efinite_map_2EFEMPTY \\ & A_27a A_27b)) V0f))) \end{aligned}$$