

thm_2Efinite__map_2ESUBMAP__REFL (TMKer9EBjzbeEyrmAHCmuYnMurPG35J2wok)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2E$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Let $ty_2Eone_2Eone : \iota$ be given. Assume the following.

$$nonempty\ ty_2Eone_2Eone \tag{1}$$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Esum_2Esum\ A0\ A1) \tag{2}$$

Let $ty_2Efinite_map_2E fmap : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Efinite_map_2E fmap\ A0\ A1) \tag{3}$$

Let $c_2Efinite_map_2E fmap_REP : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Efinite_map_2E fmap_REP\ A_27a\ A_27b \in (((ty_2Esum_2Esum\ A_27b\ ty_2Eone_2Eone)^{A_27a})(ty_2Efinite_map_2E fmap\ A_27a\ A_27b)) \tag{4}$$

Let $c_2Esum_2EOUTL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Esum_2EOUTL\ A_27a\ A_27b \in (A_27a^{(ty_2Esum_2Esum\ A_27a\ A_27b)}) \tag{5}$$

Definition 7 We define $c_2Efinite_map_2EFAPPLY$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (ty_2Efinite_map_2EFAPPLY)$

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2)))$

Let $c_2Esum_2EISL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Esum_2EISL \\ A_27a A_27b \in (2^{(ty_2Esum_2Esum A_27a A_27b)}) \end{aligned} \quad (6)$$

Definition 9 We define $c_2Efinite_map_2EFDOM$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (ty_2Efinite_map_2EFDOM)$

Definition 10 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 11 We define $c_2Efinite_map_2ESUBMAP$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (ty_2Efinite_map_2ESUBMAP)$

Assume the following.

$$True \quad (7)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in \\ A_27a.(p V0t)) \Leftrightarrow (p V0t))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow \\ True)) \end{aligned} \quad (11)$$

Theorem 1

$$\begin{aligned} \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\\ \forall V0f \in (ty_2Efinite_map_2Efmap A_27a A_27b).(p (ap (ap \\ (c_2Efinite_map_2ESUBMAP A_27a A_27b) V0f) V0f))) \end{aligned}$$