

thm_2Efinite__map_2Efmap__eq__flookup (TMPTcaja2uNmeUXSdvk66UpmJDshehLugse)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $ty_2Eone_2Eone : \iota$ be given. Assume the following.

$$nonempty\ ty_2Eone_2Eone \tag{1}$$

Definition 3 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then $(the (\lambda x.x \in A \wedge p))$ of type $\iota \Rightarrow \iota$.

Definition 4 We define c_2Eone_2Eone to be $(ap (c_2Emin_2E_40 ty_2Eone_2Eone) (\lambda V0x \in ty_2Eone_2Eone))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 6 We define $c_2Ebool_2E_F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 7 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 8 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_F))$

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Esum_2Esum\ A0\ A1) \tag{2}$$

Let $c_2Esum_2EABS_sum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Esum_2EABS_sum\ A_27a\ A_27b \in ((ty_2Esum_2Esum\ A_27a\ A_27b)^{((2^{A_27b})^{A_27a})^2}) \tag{3}$$

Definition 10 We define c_Esum_2EINR to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0e \in A_27b. (ap (c_Esum_2EABS$
Let $ty_2Eoption_2Eoption : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. nonempty A0 \Rightarrow nonempty (ty_2Eoption_2Eoption A0) \quad (4)$$

Let $c_2Eoption_2Eoption_ABS : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow c_2Eoption_2Eoption_ABS A_27a \in ((ty_2Eoption_2Eoption A_27a)^{(ty_2Esum_2Esum A_27a ty_2Eone_2Eone)}) \quad (5)$$

Definition 11 We define $c_2Eoption_2ENONE$ to be $\lambda A_27a : \iota. (ap (c_2Eoption_2Eoption_ABS A_27a) ($
Let $ty_2Efinite_map_2E fmap : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. nonempty A0 \Rightarrow \forall A1. nonempty A1 \Rightarrow nonempty (ty_2Efinite_map_2E fmap A0 A1) \quad (6)$$

Let $c_2Efinite_map_2E fmap_REP : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow c_2Efinite_map_2E fmap_REP A_27a A_27b \in (((ty_2Esum_2Esum A_27b ty_2Eone_2Eone)^{A_27a})^{(ty_2Efinite_map_2E fmap A_27a A_27b)}) \quad (7)$$

Let $c_2Esum_2EOUTL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow c_2Esum_2EOUTL A_27a A_27b \in (A_27a)^{(ty_2Esum_2Esum A_27a A_27b)} \quad (8)$$

Definition 12 We define $c_2Efinite_map_2EFAPPLY$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0f \in (ty_2Efinite_map$

Definition 13 We define c_2Esum_2EINL to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0e \in A_27a. (ap (c_2Esum_2EABS$

Definition 14 We define $c_2Eoption_2ESOME$ to be $\lambda A_27a : \iota. \lambda V0x \in A_27a. (ap (c_2Eoption_2Eoption$

Let $c_2Esum_2EISL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow c_2Esum_2EISL A_27a A_27b \in (2^{(ty_2Esum_2Esum A_27a A_27b)}) \quad (9)$$

Definition 15 We define $c_2Efinite_map_2EFDOM$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0f \in (ty_2Efinite_map$

Definition 16 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota. (\lambda V0x \in A_27a. (\lambda V1f \in (2^{A_27a}). (ap V1f V0x)))$

Definition 17 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A_27a. (\lambda V2t2 \in A_27a. ($

Definition 18 We define $c_2Efinite_map_2EFLOOKUP$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0f \in (ty_2Efinite_map$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\ \forall V0f \in (A_27b^{A_27a}).(\forall V1g \in (A_27b^{A_27a}).((V0f = \\ V1g) \Leftrightarrow (\forall V2x \in A_27a.((ap\ V0f\ V2x) = (ap\ V1g\ V2x)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\ \forall V0f1 \in (ty_2Efinite_map_2E fmap\ A_27a\ A_27b).(\forall V1f2 \in \\ (ty_2Efinite_map_2E fmap\ A_27a\ A_27b).((V0f1 = V1f2) \Leftrightarrow ((ap\ (c_2Efinite_map_2EFLOOKUP \\ A_27a\ A_27b)\ V0f1) = (ap\ (c_2Efinite_map_2EFLOOKUP\ A_27a\ A_27b) \\ V1f2)))))) \end{aligned} \quad (11)$$

Theorem 1

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\ \forall V0f1 \in (ty_2Efinite_map_2E fmap\ A_27a\ A_27b).(\forall V1f2 \in \\ (ty_2Efinite_map_2E fmap\ A_27a\ A_27b).((V0f1 = V1f2) \Leftrightarrow (\forall V2x \in \\ A_27a.((ap\ (ap\ (c_2Efinite_map_2EFLOOKUP\ A_27a\ A_27b)\ V0f1) \\ V2x) = (ap\ (ap\ (c_2Efinite_map_2EFLOOKUP\ A_27a\ A_27b)\ V1f2)\ V2x)))))) \end{aligned}$$