

thm\_2Efinite\_\_map\_2Efmap\_\_rel\_\_FEMPTY  
(TMMTcH-  
fqsAvCJkb8m2pZ5951hxBqJPKyXT9)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

Let  $ty\_2Eone\_2Eone : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Eone\_2Eone \tag{1}$$

**Definition 5** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p))$  of type  $\iota \Rightarrow \iota$ .

**Definition 6** We define  $c\_2Eone\_2Eone$  to be  $(ap (c\_2Emin\_2E\_40 ty\_2Eone\_2Eone) (\lambda V0x \in ty\_2Eone\_2Eone))$

**Definition 7** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 8** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F))$

**Definition 9** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Esum\_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Esum\_2Esum\ A0\ A1) \tag{2}$$

Let  $c\_2Esum\_2EABS\_sum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EABS\_sum\ A\_27a\ A\_27b \in ((ty\_2Esum\_2Esum\ A\_27a\ A\_27b)^{((2^{A\_27b})^{A\_27a})^2}) \tag{3}$$

**Definition 10** We define  $c\_2Esum\_2EINR$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0e \in A\_27b. (ap (c\_2Esum\_2EABS$

Let  $ty\_2Efinite\_map\_2Efmap : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Efinite\_map\_2Efmap\ A0\ A1) \quad (4)$$

Let  $c\_2Efinite\_map\_2Efmap\_ABS : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Efinite\_map\_2Efmap\_ABS\ A\_27a\ A\_27b \in ((ty\_2Efinite\_map\_2Efmap\ A\_27a\ A\_27b)^{(ty\_2Esum\_2Esum\ A\_27b\ ty\_2Eone\_2Eone)^{A\_27a}}) \quad (5)$$

**Definition 11** We define  $c\_2Efinite\_map\_2EFEMPTY$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. (ap (c\_2Efinite\_map\_2E$

Let  $c\_2Efinite\_map\_2Efmap\_REP : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Efinite\_map\_2Efmap\_REP\ A\_27a\ A\_27b \in (((ty\_2Esum\_2Esum\ A\_27b\ ty\_2Eone\_2Eone)^{A\_27a})^{(ty\_2Efinite\_map\_2Efmap\ A\_27a\ A\_27b)}) \quad (6)$$

Let  $c\_2Esum\_2EOUTL : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EOUTL\ A\_27a\ A\_27b \in (A\_27a^{(ty\_2Esum\_2Esum\ A\_27a\ A\_27b)}) \quad (7)$$

**Definition 12** We define  $c\_2Efinite\_map\_2EFAPPLY$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0f \in (ty\_2Efinite\_map$

Let  $c\_2Esum\_2EISL : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EISL\ A\_27a\ A\_27b \in (2^{(ty\_2Esum\_2Esum\ A\_27a\ A\_27b)}) \quad (8)$$

**Definition 13** We define  $c\_2Efinite\_map\_2EFDOM$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0f \in (ty\_2Efinite\_map$

**Definition 14** We define  $c\_2Ebool\_2EIN$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. (\lambda V1f \in (2^{A\_27a}). (ap\ V1f\ V0x)))$

**Definition 15** We define  $c\_2Efinite\_map\_2Efmap\_rel$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda A\_27c : \iota. \lambda V0R \in ((2^{A\_2$

**Definition 16** We define  $c\_2Epred\_set\_2EEMPTY$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. c\_2Ebool\_2EF)$ .

Assume the following.

$$True \quad (9)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (10)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p V0t))) \quad (11)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a. (p V0t)) \Leftrightarrow (p V0t))) \quad (12)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (13)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (( \\ & (p V0t) \Rightarrow False) \Leftrightarrow \neg(p V0t)))))) \end{aligned} \quad (14)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (15)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (16)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p V0t1) \Rightarrow \\ & ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2. (\forall V1x\_27 \in 2. (\forall V2y \in 2. (\forall V3y\_27 \in \\ & 2. (((p V0x) \Leftrightarrow (p V1x\_27)) \wedge ((p V1x\_27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y\_27)))) \Rightarrow \\ & (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x\_27) \Rightarrow (p V3y\_27)))))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ & (ap\ (c\_2Efinite\_map\_2EFDOM\ A\_27a\ A\_27b)\ (c\_2Efinite\_map\_2EEMPTY \\ & A\_27a\ A\_27b)) = (c\_2Epred\_set\_2EEMPTY\ A\_27a)) \end{aligned} \quad (19)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\neg(p\ (ap\ (ap\ (c\_2Ebool\_2EIN\ A\_27a)\ V0x)\ (c\_2Epred\_set\_2EEMPTY\ A\_27a)))))) \quad (20)$$

**Theorem 1**

$$\begin{aligned} & \forall A\_27a.\text{nonempty } A\_27a \Rightarrow \forall A\_27b.\text{nonempty } A\_27b \Rightarrow \forall A\_27c. \\ & \text{nonempty } A\_27c \Rightarrow (\forall V0R \in ((2^{A\_27b})^{A\_27a}).(p (ap (ap (ap ( \\ \text{c\_2Efinite\_map\_2Efm\_rel } A\_27a \ A\_27b \ A\_27c) \ V0R) \ (\text{c\_2Efinite\_map\_2EFEMPTY} \\ A\_27c \ A\_27a)) \ (\text{c\_2Efinite\_map\_2EFEMPTY } A\_27c \ A\_27b)))) \end{aligned}$$