

thm\_2Efinite\_\_map\_2Eis\_\_fmap\_\_cases  
(TMZy17zjecNtJHsUMfbbMHZuL2NV1aVkVkT)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D` to be  $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj\_o } (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define `c_2Ebool_2E_7E` to be  $(\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^2))) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x)$

**Definition 3** We define `c_2Ebool_2E_21` to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^{A\_27a}))))$

**Definition 4** We define `c_2Ebool_2E_2F` to be  $(\text{ap } (\text{c\_2Ebool\_2E\_21 } 2)) (\lambda V0t \in 2. V0t)$ .

**Definition 5** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2. \lambda Q \in 2. \text{inj\_o } (P \Rightarrow Q)$  of type  $\iota$ .

**Definition 6** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D\_3D\_3E } V0t)) (\text{c\_2Ebool\_2E\_2F})))$

**Definition 7** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2)) (\lambda V2t \in 2. V2t)))$

Let `ty_2Esum_2Esum` :  $\iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0. \text{nonempty } A0 \Rightarrow \forall A1. \text{nonempty } A1 \Rightarrow \text{nonempty } (ty\_2Esum\_2Esum \ A0 \ A1) \tag{1}$$

Let `c_2Esum_2EABS__sum` :  $\iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a. \text{nonempty } A\_27a \Rightarrow \forall A\_27b. \text{nonempty } A\_27b \Rightarrow c\_2Esum\_2EABS\_sum \ A\_27a \ A\_27b \in ((ty\_2Esum\_2Esum \ A\_27a \ A\_27b)^{((2^{A\_27b})^{A\_27a})^2}) \tag{2}$$

**Definition 8** We define `c_2Esum_2EINL` to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0e \in A\_27a. (\text{ap } (\text{c\_2Esum\_2EABS\_sum } V0e))$

**Definition 9** We define `c_2Emin_2E_40` to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. P \ x) \text{ then } (the \ (\lambda x. x \in A \wedge P \ x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define `c_2Ebool_2ECOND` to be  $\lambda A\_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A\_27a. (\lambda V2t2 \in A\_27a. (V2t2 \in A\_27a))))$

**Definition 11** We define `c_2Ebool_2E_3F` to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (\text{ap } V0P \ (\text{ap } (\text{c\_2Emin\_2E\_40 } V0P))))$

Let  $ty\_2Eone\_2Eone : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Eone\_2Eone \quad (3)$$

**Definition 12** We define  $c\_2Eone\_2Eone$  to be  $(ap\ (c\_2Emin\_2E\_40\ ty\_2Eone\_2Eone)\ (\lambda V0x \in ty\_2Eone\_2Eone))$

**Definition 13** We define  $c\_2Esum\_2EINR$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27b.(ap\ (c\_2Esum\_2EABS$

**Definition 14** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 15** We define  $c\_2Efinite\_map\_2Eis\_fmap$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.(\lambda V0a0 \in ((ty\_2Esum\_2E$

Assume the following.

$$True \quad (4)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (5)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (6)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1y \in 2.(\forall V2z \in 2.(\forall V3w \in 2.(((p\ V0x) \Rightarrow (p\ V1y)) \wedge ((p\ V2z) \Rightarrow (p\ V3w))) \Rightarrow (((p\ V0x) \wedge (p\ V2z)) \Rightarrow ((p\ V1y) \wedge (p\ V3w)))))) \quad (7)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1y \in 2.(\forall V2z \in 2.(\forall V3w \in 2.(((p\ V0x) \Rightarrow (p\ V1y)) \wedge ((p\ V2z) \Rightarrow (p\ V3w))) \Rightarrow (((p\ V0x) \vee (p\ V2z)) \Rightarrow ((p\ V1y) \vee (p\ V3w)))))) \quad (8)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}).(\forall V1Q \in (2^{A\_27a}).((\forall V2x \in A\_27a.((p\ (ap\ V0P\ V2x)) \Rightarrow (p\ (ap\ V1Q\ V2x)))) \Rightarrow ((\exists V3x \in A\_27a.(p\ (ap\ V0P\ V3x))) \Rightarrow (\exists V4x \in A\_27a.(p\ (ap\ V1Q\ V4x)))))) \quad (9)$$

**Theorem 1**

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ & \quad \forall V0a0 \in ((ty\_2Esum\_2Esum\ A\_27b\ ty\_2Eone\_2Eone)^{A\_27a}). \\ & \quad ((p\ (ap\ (c\_2Efinite\_map\_2Eis\_fmap\ A\_27a\ A\_27b)\ V0a0)) \Leftrightarrow ((V0a0 = \\ & (\lambda V1a \in A\_27a.(ap\ (c\_2Esum\_2EINR\ A\_27b\ ty\_2Eone\_2Eone)\ c\_2Eone\_2Eone))) \vee \\ & \quad (\exists V2f \in ((ty\_2Esum\_2Esum\ A\_27b\ ty\_2Eone\_2Eone)^{A\_27a}). \\ & \quad (\exists V3a \in A\_27a.(\exists V4b \in A\_27b.((V0a0 = (\lambda V5x \in A\_27a. \\ & (ap\ (ap\ (c\_2Ebool\_2ECOND\ (ty\_2Esum\_2Esum\ A\_27b\ ty\_2Eone\_2Eone)) \\ & (ap\ (ap\ (c\_2Emin\_2E\_3D\ A\_27a)\ V5x)\ V3a))\ (ap\ (c\_2Esum\_2EINL\ A\_27b \\ & ty\_2Eone\_2Eone)\ V4b))\ (ap\ V2f\ V5x)))) \wedge (p\ (ap\ (c\_2Efinite\_map\_2Eis\_fmap \\ & \quad A\_27a\ A\_27b)\ V2f)))))) \end{aligned}$$