

thm_2Egcd_2ELCM_0 (TMSR- BxS4Sp1BtwuVKDCwRbFWqc352n9J6jC)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Egcd_2Egcd : \iota$ be given. Assume the following.

$$c_2Egcd_2Egcd \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Earithmetic_2E_2A : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2A \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{3}$$

Let $c_2Earithmetic_2EDIV : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EDIV \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{5}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{6}$$

Definition 8 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 9 We define $c_2Ebool_2E5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E21\ 2)\ (\lambda V2t \in 2.$

Definition 10 We define c_2Emin_2E40 to be $\lambda A.\lambda P \in 2^A$. **if** $(\exists x \in A.p\ (ap\ P\ x))$ **then** $(the\ (\lambda x.x \in A \wedge P\ x))$ of type $\iota \Rightarrow \iota$.

Definition 11 We define c_2Ebool_2ECOND to be $\lambda A.\lambda a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A.\lambda V2t2 \in A.\lambda V2t1 \in A.$

Definition 12 We define c_2Egcd_2Elcm to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.(ap\ (ap\ c_2Egcd_2Elcm\ V0m\ V1n))$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum.((ap\ (ap\ c_2Earithmetic_2E_2A\ V0m)\ c_2Enum_2E0) = c_2Enum_2E0)) \quad (7)$$

Assume the following.

$$(ap\ (ap\ c_2Earithmetic_2E_2A\ V0m)\ V1n) = (ap\ (ap\ c_2Earithmetic_2E_2A\ V1n)\ V0m)) \quad (8)$$

Assume the following.

$$True \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.(((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \quad (10)$$

Assume the following.

$$\forall A.\lambda a.nonempty\ A.\lambda a \Rightarrow (\forall V0x \in A.\lambda a.((V0x = V0x) \Leftrightarrow True)) \quad (11)$$

Assume the following.

$$\forall A.\lambda a.nonempty\ A.\lambda a \Rightarrow (\forall V0x \in A.\lambda a.(\forall V1y \in A.\lambda a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (12)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p\ V0t1) \Rightarrow ((p\ V1t2) \Rightarrow (p\ V2t3))) \Leftrightarrow (((p\ V0t1) \wedge (p\ V1t2)) \Rightarrow (p\ V2t3)))))) \quad (13)$$

Assume the following.

$$\begin{aligned} & \forall A.\lambda a.nonempty\ A.\lambda a \Rightarrow (\forall V0P \in 2.(\forall V1Q \in 2. \\ & (\forall V2x \in A.\lambda a.(\forall V3x_27 \in A.\lambda a.(\forall V4y \in A.\lambda a. \\ & (\forall V5y_27 \in A.\lambda a.(((p\ V0P) \Leftrightarrow (p\ V1Q)) \wedge (((p\ V1Q) \Rightarrow (V2x = V3x_27)) \wedge \\ & ((\neg(p\ V1Q)) \Rightarrow (V4y = V5y_27)))))) \Rightarrow ((ap\ (ap\ (ap\ (c_2Ebool_2ECOND\ A.\lambda a)\ V0P)\ V2x)\ V4y) = (ap\ (ap\ (ap\ (c_2Ebool_2ECOND\ A.\lambda a)\ V1Q)\ V3x_27)\ V5y_27)))))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow & ((\forall V0t1 \in A_27a. (\forall V1t2 \in \\ A_27a. & ((ap\ (ap\ (ap\ (c_2Ebool_2ECOND\ A_27a)\ c_2Ebool_2ET)\ V0t1) \\ & V1t2) = V0t1))) \wedge (\forall V2t1 \in A_27a. (\forall V3t2 \in A_27a. ((ap \\ & (ap\ (ap\ (c_2Ebool_2ECOND\ A_27a)\ c_2Ebool_2EF)\ V2t1)\ V3t2) = V3t2)))) \end{aligned} \quad (15)$$

Assume the following.

$$(\forall V0a \in ty_2Enum_2Enum. ((ap\ (ap\ c_2Egcd_2Egcd\ V0a)\ c_2Enum_2E0) = V0a)) \quad (16)$$

Assume the following.

$$(\forall V0a \in ty_2Enum_2Enum. ((ap\ (ap\ c_2Egcd_2Egcd\ c_2Enum_2E0)\ V0a) = V0a)) \quad (17)$$

Theorem 1

$$(\forall V0x \in ty_2Enum_2Enum. (((ap\ (ap\ c_2Egcd_2Elcm\ c_2Enum_2E0)\ V0x) = c_2Enum_2E0) \wedge ((ap\ (ap\ c_2Egcd_2Elcm\ V0x)\ c_2Enum_2E0) = c_2Enum_2E0)))$$