

thm\_2Ehreal\_2EHRAT\_\_DOWN  
(TMNKWkqufgStbtdrtYBwnZ3rizpjPTRb7e8)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Ehrat\_2Ehrat : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehrat\_2Ehrat \tag{3}$$

Let  $c\_2Ehrat\_2Ehrat\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Ehrat\_2Ehrat\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)})_{ty\_2Ehrat\_2Ehrat}) \tag{4}$$

**Definition 1** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A$ .if  $(\exists x \in A.p\ (ap\ P\ x))$  **then** (the  $(\lambda x.x \in A \wedge p\ x)$  of type  $\iota \Rightarrow \iota$ ).

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o\ (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E\_2ET$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

**Definition 5** We define  $c\_2Ehrat\_2Ehrat\_REP$  to be  $\lambda V0a \in ty\_2Ehrat\_2Ehrat.(ap\ (c\_2Emin\_2E\_40\ (ty\_2Ehrat\_2Ehrat\ a)))$

Let  $c\_2Ehrat\_2Etrac\_add : \iota$  be given. Assume the following.

$$c\_2Ehrat\_2Etrac\_add \in (((ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum)})_{c\_2Ehrat\_2Etrac\_add}) \tag{5}$$



Let  $c\_2Ehrat\_2Etrat\_mul : \iota$  be given. Assume the following.

$$c\_2Ehrat\_2Etrat\_mul \in (((ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum)})) \quad (12)$$

**Definition 16** We define  $c\_2Ehrat\_2Ehtrat\_mul$  to be  $\lambda V0T1 \in ty\_2Ehtrat\_2Ehtrat.\lambda V1T2 \in ty\_2Ehtrat\_2Ehtrat$

**Definition 17** We define  $c\_2Ehreal\_2Ehtrat\_lt$  to be  $\lambda V0x \in ty\_2Ehtrat\_2Ehtrat.\lambda V1y \in ty\_2Ehtrat\_2Ehtrat$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (13)$$

Assume the following.

$$(\forall V0h \in ty\_2Ehtrat\_2Ehtrat. ((ap\ (ap\ c\_2Ehtrat\_2Ehtrat\_mul\ c\_2Ehtrat\_2Ehtrat\_1)\ V0h) = V0h)) \quad (14)$$

Assume the following.

$$(\forall V0x \in ty\_2Ehtrat\_2Ehtrat. (\forall V1y \in ty\_2Ehtrat\_2Ehtrat. ((p\ (ap\ (ap\ c\_2Ehreal\_2Ehtrat\_lt\ (ap\ (ap\ c\_2Ehtrat\_2Ehtrat\_mul\ V0x)\ V1y))\ V0x)) \Leftrightarrow (p\ (ap\ (ap\ c\_2Ehreal\_2Ehtrat\_lt\ V1y)\ c\_2Ehtrat\_2Ehtrat\_1)))))) \quad (15)$$

Assume the following.

$$(\forall V0x \in ty\_2Ehtrat\_2Ehtrat. (\forall V1y \in ty\_2Ehtrat\_2Ehtrat. ((p\ (ap\ (ap\ c\_2Ehreal\_2Ehtrat\_lt\ (ap\ (ap\ c\_2Ehtrat\_2Ehtrat\_mul\ V0x)\ (ap\ c\_2Ehtrat\_2Ehtrat\_inv\ V1y)))\ c\_2Ehtrat\_2Ehtrat\_1)) \Leftrightarrow (p\ (ap\ (ap\ c\_2Ehreal\_2Ehtrat\_lt\ V0x)\ V1y)))))) \quad (16)$$

**Theorem 1**

$$(\forall V0x \in ty\_2Ehtrat\_2Ehtrat. (\exists V1y \in ty\_2Ehtrat\_2Ehtrat. (p\ (ap\ (ap\ c\_2Ehreal\_2Ehtrat\_lt\ V1y)\ V0x))))$$