

# thm\_2Ehreal\_2EHRAT\_\_EQ\_\_LADD (TMcnodcn9bZfBczP54pjNcTKRq2eqzDNLGJ)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

**Definition 7** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Ehrat\_2Ehrat : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Ehrat\_2Ehrat \tag{3}$$

Let  $c\_2Ehrat\_2Ehrat\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Ehrat\_2Ehrat\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)})ty\_2Ehrat\_2Ehrat) \tag{4}$$

**Definition 8** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 9** We define  $c\_2Ehrat\_2Ehtrat\_REP$  to be  $\lambda V0a \in ty\_2Ehtrat\_2Ehtrat.(ap (c\_2Emin\_2E\_40 (ty\_2E$

Let  $c\_2Ehtrat\_2Etrat\_add : \iota$  be given. Assume the following.

$$c\_2Ehtrat\_2Etrat\_add \in (((ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum)})) \quad (5)$$

Let  $c\_2Ehtrat\_2Etrat\_eq : \iota$  be given. Assume the following.

$$c\_2Ehtrat\_2Etrat\_eq \in ((2^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)})^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum)}) \quad (6)$$

Let  $c\_2Ehtrat\_2Ehtrat\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Ehtrat\_2Ehtrat\_ABS\_CLASS \in (ty\_2Ehtrat\_2Ehtrat^{(2^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)})}) \quad (7)$$

**Definition 10** We define  $c\_2Ehtrat\_2Ehtrat\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2$

**Definition 11** We define  $c\_2Ehtrat\_2Ehtrat\_add$  to be  $\lambda V0T1 \in ty\_2Ehtrat\_2Ehtrat.\lambda V1T2 \in ty\_2Ehtrat\_2E$

**Definition 12** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40$

**Definition 13** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \quad (10)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow \neg (p V0t)))))) \quad (11)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (12)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (13)$$

Assume the following.

$$\begin{aligned}
& (\forall V0h \in ty\_2Ehrt\_2Ehrt. (\forall V1i \in ty\_2Ehrt\_2Ehrt. \\
& (\forall V2j \in ty\_2Ehrt\_2Ehrt. ((ap (ap c\_2Ehrt\_2Ehrt\_add \\
V0h) (ap (ap c\_2Ehrt\_2Ehrt\_add V1i) V2j)) = (ap (ap c\_2Ehrt\_2Ehrt\_add \\
& (ap (ap c\_2Ehrt\_2Ehrt\_add V0h) V1i)) V2j))))))
\end{aligned} \tag{14}$$

Assume the following.

$$\begin{aligned}
& (\forall V0h \in ty\_2Ehrt\_2Ehrt. (\forall V1i \in ty\_2Ehrt\_2Ehrt. \\
& (\neg((ap (ap c\_2Ehrt\_2Ehrt\_add V0h) V1i) = V0h))))
\end{aligned} \tag{15}$$

Assume the following.

$$\begin{aligned}
& (\forall V0h \in ty\_2Ehrt\_2Ehrt. (\forall V1i \in ty\_2Ehrt\_2Ehrt. \\
& ((V0h = V1i) \vee ((\exists V2d \in ty\_2Ehrt\_2Ehrt. (V0h = (ap (ap c\_2Ehrt\_2Ehrt\_add \\
V1i) V2d))) \vee (\exists V3d \in ty\_2Ehrt\_2Ehrt. (V1i = (ap (ap c\_2Ehrt\_2Ehrt\_add \\
& V0h) V3d)))))))
\end{aligned} \tag{16}$$

**Theorem 1**

$$\begin{aligned}
& (\forall V0x \in ty\_2Ehrt\_2Ehrt. (\forall V1y \in ty\_2Ehrt\_2Ehrt. \\
& (\forall V2z \in ty\_2Ehrt\_2Ehrt. (((ap (ap c\_2Ehrt\_2Ehrt\_add \\
V0x) V1y) = (ap (ap c\_2Ehrt\_2Ehrt\_add V0x) V2z)) \Leftrightarrow (V1y = V2z))))))
\end{aligned}$$