

thm_2Ehreal_2EHRAT__LT__GT
(TMT8uHGN98tAARXGbd8FueUFEcyn8HiXBYc)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define `c_2Emin_2E_3D` to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define `c_2Ebool_2ET` to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define `c_2Ebool_2E_21` to be $\lambda A_{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 5 We define `c_2Ebool_2EF` to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 6 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 7 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Let `ty_2Enum_2Enum` : ι be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let `ty_2Epair_2Eprod` : $\iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let `ty_2Ehrat_2Ehrat` : ι be given. Assume the following.

$$nonempty\ ty_2Ehrat_2Ehrat \tag{3}$$

Let `c_2Ehrat_2Ehrat__REP__CLASS` : ι be given. Assume the following.

$$c_2Ehrat_2Ehrat_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})ty_2Ehrat_2Ehrat) \tag{4}$$

Definition 8 We define `c_2Emin_2E_40` to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 9 We define $c_2Eh_rat_2Eh_rat_REP$ to be $\lambda V0a \in ty_2Eh_rat_2Eh_rat.(ap (c_2Emin_2E_40 (ty_2E$

Let $c_2Eh_rat_2Eh_rat_add : \iota$ be given. Assume the following.

$$c_2Eh_rat_2Eh_rat_add \in (((ty_2Epair_2Eprod ty_2Enum_2Enum ty_2Enum_2Enum)^{(ty_2Epair_2Eprod ty_2Enum_2Enum)})) \quad (5)$$

Let $c_2Eh_rat_2Eh_rat_eq : \iota$ be given. Assume the following.

$$c_2Eh_rat_2Eh_rat_eq \in ((2^{(ty_2Epair_2Eprod ty_2Enum_2Enum ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod ty_2Enum_2Enum)}) \quad (6)$$

Let $c_2Eh_rat_2Eh_rat_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Eh_rat_2Eh_rat_ABS_CLASS \in (ty_2Eh_rat_2Eh_rat^{(2^{(ty_2Epair_2Eprod ty_2Enum_2Enum ty_2Enum_2Enum)})}) \quad (7)$$

Definition 10 We define $c_2Eh_rat_2Eh_rat_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod ty_2Enum_2Enum ty_2$

Definition 11 We define $c_2Eh_rat_2Eh_rat_add$ to be $\lambda V0T1 \in ty_2Eh_rat_2Eh_rat.\lambda V1T2 \in ty_2Eh_rat_2E$

Definition 12 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40$

Definition 13 We define $c_2Eh_real_2Eh_rat_lt$ to be $\lambda V0x \in ty_2Eh_rat_2Eh_rat.\lambda V1y \in ty_2Eh_rat_2Eh_rat$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \quad (10)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\exists V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (11)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (12)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (13)$$

Assume the following.

$$\begin{aligned}
& (\forall V0h \in ty_2Ehrtat_2Ehrtat. (\forall V1i \in ty_2Ehrtat_2Ehrtat. \\
& (\forall V2j \in ty_2Ehrtat_2Ehrtat. ((ap (ap c_2Ehrtat_2Ehrtat_add \\
V0h) (ap (ap c_2Ehrtat_2Ehrtat_add V1i) V2j)) = (ap (ap c_2Ehrtat_2Ehrtat_add \\
& (ap (ap c_2Ehrtat_2Ehrtat_add V0h) V1i)) V2j))))))
\end{aligned} \tag{14}$$

Assume the following.

$$\begin{aligned}
& (\forall V0h \in ty_2Ehrtat_2Ehrtat. (\forall V1i \in ty_2Ehrtat_2Ehrtat. \\
& (\neg((ap (ap c_2Ehrtat_2Ehrtat_add V0h) V1i) = V0h))))
\end{aligned} \tag{15}$$

Theorem 1

$$\begin{aligned}
& (\forall V0x \in ty_2Ehrtat_2Ehrtat. (\forall V1y \in ty_2Ehrtat_2Ehrtat. \\
& ((p (ap (ap c_2Ehrtat_2Ehrtat_lt V0x) V1y)) \Rightarrow (\neg(p (ap (ap c_2Ehrtat_2Ehrtat_lt \\
& V1y) V0x)))))
\end{aligned}$$