

thm_2Eind_type_2EINJA_INJ
(TMV2u2Xaaajrm4AKtZcMHNXqz82HgRNPpbka)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Definition 8 We define $c_2Eind_type_2EINJA$ to be $\lambda A_27a : \iota.\lambda V0a \in A_27a.(\lambda V1n \in ty_2Enum_2Enum$

Assume the following.

$$True \tag{2}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{3}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \tag{4}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\
& True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\
& (p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t))))))
\end{aligned} \tag{5}$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0x \in A.27a.((V0x = V0x) \Leftrightarrow True)) \tag{6}$$

Assume the following.

$$\begin{aligned}
& \forall A.27a.nonempty A.27a \Rightarrow \forall A.27b.nonempty A.27b \Rightarrow (\\
& \forall V0f \in (A.27b^{A.27a}).(\forall V1g \in (A.27b^{A.27a}).((V0f = \\
& V1g) \Leftrightarrow (\forall V2x \in A.27a.((ap V0f V2x) = (ap V1g V2x))))))
\end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\
& (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(\\
& p V0t))))))
\end{aligned} \tag{8}$$

Theorem 1

$$\begin{aligned}
& \forall A.27a.nonempty A.27a \Rightarrow (\forall V0a1 \in A.27a.(\forall V1a2 \in \\
& A.27a.(((ap (c.2Eind_type_2EINJA A.27a) V0a1) = (ap (c.2Eind_type_2EINJA \\
& A.27a) V1a2)) \Leftrightarrow (V0a1 = V1a2))))
\end{aligned}$$