

thm_2Eint__arith_2EINT__NUM__EVEN (TMF-
fLLcUDBXH8jRm8WVJuHfRFxE8UW6bEZw)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Earithmetic_2EEVEN : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EEVEN \in (2^{ty_2Enum_2Enum}) \tag{2}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in omega \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{omega}) \tag{4}$$

Definition 7 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Definition 8 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (5)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (6)$$

Definition 9 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ ($

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (7)$$

Definition 10 We define $c_2Earithmetic_2EBIT2$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap\ (ap\ c_2Earithmetic_2E_2B\ ($

Definition 11 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \quad (8)$$

Let $c_2Einteger_2Eint_of_num : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_of_num \in (ty_2Einteger_2Eint^{ty_2Enum_2Enum}) \quad (9)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (10)$$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \quad (11)$$

Definition 12 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p\ (ap\ P\ x))$ then $(the\ (\lambda x.x \in A \wedge P\ x))$ of type $\iota \Rightarrow \iota$.

Definition 13 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint.(ap\ (c_2Emin_2E_40\ ($

Let $c_2Einteger_2Etint_mul : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_mul \in (((ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)}) \quad (12)$$

Let $c_2Einteger_2Etint_eq : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)}) \quad (13)$$

Let $c_2Einteger_2Eint_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_ABS_CLASS \in (ty_2Einteger_2Eint^{(2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})}) \quad (14)$$

Definition 14 We define `c_Integer.Eint_ABS` to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2$

Definition 15 We define `c_Integer.Eint_mul` to be $\lambda V0T1 \in ty_2Einteger_2Eint.\lambda V1T2 \in ty_2Eintege$

Definition 16 We define `c_Ebool_2E_3F` to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ V0P\ (ap\ (c_2Emin_2E_40$

Definition 17 We define `c_Integer.Eint_divides` to be $\lambda V0p \in ty_2Einteger_2Eint.\lambda V1q \in ty_2Eintege$

Definition 18 We define `c_Ebool_2E_7E` to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2E$

Let `c_Earithmetic_2EODD` : ι be given. Assume the following.

$$c_2Earithmetic_2EODD \in (2^{ty_2Enum_2Enum}) \quad (15)$$

Assume the following.

$$\begin{aligned} (\forall V0n \in ty_2Enum_2Enum.((p\ (ap\ c_2Earithmetic_2EEVEN\ V0n)) \Leftrightarrow \\ (\neg(p\ (ap\ c_2Earithmetic_2EODD\ V0n)))))) \end{aligned} \quad (16)$$

Assume the following.

$$True \quad (17)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (18)$$

Assume the following.

$$\begin{aligned} ((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge \\ ((\neg False) \Leftrightarrow True))) \end{aligned} \quad (19)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (20)$$

Assume the following.

$$\begin{aligned} (\forall V0n \in ty_2Enum_2Enum.((p\ (ap\ c_2Earithmetic_2EODD\ V0n)) \Leftrightarrow \\ (\neg(p\ (ap\ (ap\ c_2Einteger_2Eint_divides\ (ap\ c_2Einteger_2Eint_of_num \\ (ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT2\ c_2Earithmetic_2EZERO)))) \\ (ap\ c_2Einteger_2Eint_of_num\ V0n)))))) \end{aligned} \quad (21)$$

Theorem 1

$$\begin{aligned} (\forall V0n \in ty_2Enum_2Enum.((p\ (ap\ c_2Earithmetic_2EEVEN\ V0n)) \Leftrightarrow \\ (p\ (ap\ (ap\ c_2Einteger_2Eint_divides\ (ap\ c_2Einteger_2Eint_of_num \\ (ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT2\ c_2Earithmetic_2EZERO)))) \\ (ap\ c_2Einteger_2Eint_of_num\ V0n)))))) \end{aligned}$$