

thm_2Eint_arith_2Elim_neg_ones (TMHTJv-
Tykbs84Gnybotu9QPKCtpwkDFK5sU)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V1x \in 2.V1x))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define $c_2Earithmic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmic_2EBIT1) n)$

Definition 8 We define $c_2Earithmic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \quad (7)$$

Let $c_2Einteger_2Eint_of_num : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_of_num \in (ty_2Einteger_2Eint^{ty_2Enum_2Enum}) \quad (8)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (9)$$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \quad (10)$$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap\ P\ x)) \text{ then } (the\ (\lambda x.x \in A \wedge p\ x)) \text{ of type } \iota \Rightarrow \iota$.

Definition 10 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint.(ap\ (c_2Emin_2E_40\ a))$

Let $c_2Einteger_2Eint_neg : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_neg \in ((ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)^{ty_2Einteger_2Eint}) \quad (11)$$

Let $c_2Einteger_2Eint_eq : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \quad (12)$$

Let $c_2Einteger_2Eint_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_ABS_CLASS \in (ty_2Einteger_2Eint^{(2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})}) \quad (13)$$

Definition 11 We define $c_2Einteger_2Eint_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)$

Definition 12 We define $c_2Einteger_2Eint_neg$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.(ap\ c_2Einteger_2Eint_neg)$

Let $c_2Einteger_2Eint_add : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_add \in (((ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)^{ty_2Einteger_2Eint})^{ty_2Einteger_2Eint}) \quad (14)$$

Definition 13 We define $c_Einteger_2Eint_add$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.\lambda V1T2 \in ty_2Einteger_2Eint$

Assume the following.

$$True \tag{15}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t) \Leftrightarrow (p\ V0t)))) \tag{16}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \tag{17}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{18}$$

Assume the following.

$$\begin{aligned} & (\forall V0z \in ty_2Einteger_2Eint.(\forall V1y \in ty_2Einteger_2Eint. \\ & (\forall V2x \in ty_2Einteger_2Eint.((ap\ (ap\ c_2Einteger_2Eint_add \\ & V2x)\ (ap\ (ap\ c_2Einteger_2Eint_add\ V1y)\ V0z)) = (ap\ (ap\ c_2Einteger_2Eint_add \\ & (ap\ (ap\ c_2Einteger_2Eint_add\ V2x)\ V1y))\ V0z)))))) \end{aligned} \tag{19}$$

Assume the following.

$$(\forall V0x \in ty_2Einteger_2Eint.((ap\ (ap\ c_2Einteger_2Eint_add\ V0x)\ (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0)) = V0x)) \tag{20}$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty_2Einteger_2Eint.((ap\ (ap\ c_2Einteger_2Eint_add \\ & (ap\ c_2Einteger_2Eint_neg\ V0x))\ V0x) = (ap\ c_2Einteger_2Eint_of_num \\ & c_2Enum_2E0))) \end{aligned} \tag{21}$$

Theorem 1

$$\begin{aligned} & (\forall V0x \in ty_2Einteger_2Eint.((ap\ (ap\ c_2Einteger_2Eint_add \\ & (ap\ (ap\ c_2Einteger_2Eint_add\ V0x)\ (ap\ c_2Einteger_2Eint_neg \\ & (ap\ c_2Einteger_2Eint_of_num\ (ap\ c_2Earithmetic_2ENUMERAL \\ & (ap\ c_2Earithmetic_2EBIT1\ c_2Earithmetic_2EZERO))))))\ (ap\ c_2Einteger_2Eint_of_num \\ & (ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT1\ c_2Earithmetic_2EZERO)))) = \\ & V0x)) \end{aligned}$$