

thm_2Eint__arith_2Eq__context__rwt2
(TMZrGkRVsCwEKMKKT8YKudHnm8hJyr5errn2)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow P \Rightarrow Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \tag{3}$$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})_{ty_2Einteger_2Eint}) \tag{4}$$

Definition 4 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap\ P\ x)) \mathbf{then} (the (\lambda x.x \in A \wedge p))$ of type $\iota \Rightarrow \iota$.

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda 27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})))$

Definition 6 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint.(ap (c_2Emin_2E_40 (ty_2Einteger_2Eint_REP_CLASS (2^{V0a})))$

Definition 14 We define $c_Einteger_2Eint_neg$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.(ap\ c_2Einteger_2Eint$.

Assume the following.

$$True \quad (13)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (14)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty_2Einteger_2Eint.(\forall V1y \in ty_2Einteger_2Eint. \\ & ((ap\ c_2Einteger_2Eint_neg\ (ap\ (ap\ c_2Einteger_2Eint_add\ V0x) \\ & V1y)) = (ap\ (ap\ c_2Einteger_2Eint_add\ (ap\ c_2Einteger_2Eint_neg \\ & V0x))\ (ap\ c_2Einteger_2Eint_neg\ V1y)))))) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty_2Einteger_2Eint.(\forall V1y \in ty_2Einteger_2Eint. \\ & (\forall V2z \in ty_2Einteger_2Eint.((p\ (ap\ (ap\ c_2Einteger_2Eint_le \\ & (ap\ (ap\ c_2Einteger_2Eint_add\ V0x)\ V1y))\ (ap\ (ap\ c_2Einteger_2Eint_add \\ & V0x)\ V2z))) \Leftrightarrow (p\ (ap\ (ap\ c_2Einteger_2Eint_le\ V1y)\ V2z)))))) \end{aligned} \quad (16)$$

Assume the following.

$$((ap\ c_2Einteger_2Eint_neg\ (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0)) = (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0)) \quad (17)$$

Theorem 1

$$\begin{aligned} & (\forall V0c \in ty_2Einteger_2Eint.(\forall V1x \in ty_2Einteger_2Eint. \\ & (\forall V2y \in ty_2Einteger_2Eint.(((ap\ c_2Einteger_2Eint_of_num \\ & c_2Enum_2E0) = (ap\ (ap\ c_2Einteger_2Eint_add\ V0c)\ V1x)) \Rightarrow ((p\ (\\ & ap\ (ap\ c_2Einteger_2Eint_le\ (ap\ c_2Einteger_2Eint_of_num \\ & c_2Enum_2E0))\ (ap\ (ap\ c_2Einteger_2Eint_add\ (ap\ c_2Einteger_2Eint_neg \\ & V0c)\ V2y))) \Leftrightarrow (p\ (ap\ (ap\ c_2Einteger_2Eint_le\ (ap\ c_2Einteger_2Eint_neg \\ & V1x))\ V2y)))))) \end{aligned}$$