

# tm2Eint\_arith2Elt\_move\_all\_right (TMUkR-CBZUBXpHMo27yowqUyhJUPkjqh87AU)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 3** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $ty\_2Einteger\_2Eint : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Einteger\_2Eint \tag{4}$$

Let  $c\_2Einteger\_2Eint\_of\_num : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_of\_num \in (ty\_2Einteger\_2Eint^{ty\_2Enum\_2Enum}) \tag{5}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{6}$$

Let  $c\_2Einteger\_2Eint\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)})\ ty\_2Einteger\_2Eint) \tag{7}$$

**Definition 4** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.$ **if**  $(\exists x \in A.p (ap P x))$  **then** *(the*  $(\lambda x.x \in A \wedge p$   
of type  $\iota \Rightarrow \iota$ .

**Definition 5** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}$

**Definition 6** We define  $c\_2Einteger\_2Eint\_REP$  to be  $\lambda V0a \in ty\_2Einteger\_2Eint.(ap (c\_2Emin\_2E\_40 (ty$

Let  $c\_2Einteger\_2Eint\_neg : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_neg \in ((ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum) (ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)) \quad (8)$$

Let  $c\_2Einteger\_2Eint\_eq : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_eq \in ((2^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)}) (ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)) \quad (9)$$

Let  $c\_2Einteger\_2Eint\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_ABS\_CLASS \in (ty\_2Einteger\_2Eint)^{(2^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)})} \quad (10)$$

**Definition 7** We define  $c\_2Einteger\_2Eint\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)$

**Definition 8** We define  $c\_2Einteger\_2Eint\_neg$  to be  $\lambda V0T1 \in ty\_2Einteger\_2Eint.(ap c\_2Einteger\_2Eint\_neg$

Let  $c\_2Einteger\_2Eint\_add : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_add \in (((ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum) (ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)) (ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)) \quad (11)$$

**Definition 9** We define  $c\_2Einteger\_2Eint\_add$  to be  $\lambda V0T1 \in ty\_2Einteger\_2Eint.\lambda V1T2 \in ty\_2Einteger\_2Eint$

Let  $c\_2Einteger\_2Eint\_lt : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_lt \in ((2^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)}) (ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)) \quad (12)$$

**Definition 10** We define  $c\_2Einteger\_2Eint\_lt$  to be  $\lambda V0T1 \in ty\_2Einteger\_2Eint.\lambda V1T2 \in ty\_2Einteger\_2Eint$

Assume the following.

$$True \quad (13)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (14)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (15)$$

Assume the following.

$$(\forall V0x \in ty\_2Einteger\_2Eint.((ap (ap c\_2Einteger\_2Eint\_add (ap c\_2Einteger\_2Eint\_of\_num c\_2Enum\_2E0)) V0x) = V0x)) \quad (16)$$

Assume the following.

$$(\forall V0x \in ty\_2Einteger\_2Eint.(\forall V1y \in ty\_2Einteger\_2Eint. (\forall V2z \in ty\_2Einteger\_2Eint.((p (ap (ap c\_2Einteger\_2Eint\_lt V1y) (ap (ap c\_2Einteger\_2Eint\_add V0x) (ap c\_2Einteger\_2Eint\_neg V2z)))) \Leftrightarrow (p (ap (ap c\_2Einteger\_2Eint\_lt (ap (ap c\_2Einteger\_2Eint\_add V1y) V2z)) V0x)))))) \quad (17)$$

**Theorem 1**

$$(\forall V0x \in ty\_2Einteger\_2Eint.(\forall V1y \in ty\_2Einteger\_2Eint. ((p (ap (ap c\_2Einteger\_2Eint\_lt V0x) V1y)) \Leftrightarrow (p (ap (ap c\_2Einteger\_2Eint\_lt (ap c\_2Einteger\_2Eint\_of\_num c\_2Enum\_2E0)) (ap (ap c\_2Einteger\_2Eint\_add V1y) (ap c\_2Einteger\_2Eint\_neg V0x))))))))$$