

thm_2Einteger_2EINT__EQ__IMP__LE (TMKZvRN8R87v9coHrR7Cvrc2iH2SgqUHbJX)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow P \Rightarrow Q)$ of type ι .

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \tag{3}$$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \tag{4}$$

Definition 3 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p)$ of type $\iota \Rightarrow \iota$).

Definition 4 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 6 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint.(ap (c_2Emin_2E_40 (ty$

Let $c_2Einteger_2Etint_lt : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \tag{5}$$

Definition 7 We define $c_Einteger_Eint_lt$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.\lambda V1T2 \in ty_2Einteger_2Eint.$

Definition 8 We define c_Ebool_EF to be $(ap (c_Ebool_E21) 2) (\lambda V0t \in 2.V0t)$.

Definition 9 We define c_Ebool_E7E to be $(\lambda V0t \in 2.(ap (ap c_Emin_E3D_3D_3E V0t) c_Ebool_EF))$

Definition 10 We define $c_Einteger_Eint_le$ to be $\lambda V0x \in ty_2Einteger_2Eint.\lambda V1y \in ty_2Einteger_2Eint.$

Assume the following.

$$(\forall V0x \in ty_2Einteger_2Eint.(p (ap (ap c_Einteger_Eint_le V0x) V0x))) \quad (6)$$

Theorem 1

$$(\forall V0x \in ty_2Einteger_2Eint.(\forall V1y \in ty_2Einteger_2Eint.((V0x = V1y) \Rightarrow (p (ap (ap c_Einteger_Eint_le V0x) V1y)))))$$