

# thm\_2Einteger\_2EINT\_EQ\_LMUL2 (TMXcJhLPTaw3aAtZZtXHRDHV1adVs9jZize)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 8** We define  $c\_2Enum\_2E0$  to be  $(ap c\_2Enum\_2EABS\_num c\_2Enum\_2EZERO\_REP)$ .

Let  $ty\_2Einteger\_2Eint : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Einteger\_2Eint \tag{4}$$

Let  $c\_2Einteger\_2Eint\_of\_num : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_of\_num \in (ty\_2Einteger\_2Eint^{ty\_2Enum\_2Enum}) \tag{5}$$

**Definition 9** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.))$   
 Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Epair\_2Eprod A0 A1) \quad (6)$$

Let  $c\_2Einteger\_2Eint\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)})^{ty\_2Einteger\_2Eint\_REP\_CLASS}) \quad (7)$$

**Definition 10** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.$ if  $(\exists x \in A.p (ap P x))$  then  $(the (\lambda x.x \in A \wedge P x))$   
 of type  $\iota \Rightarrow \iota$ .

**Definition 11** We define  $c\_2Einteger\_2Eint\_REP$  to be  $\lambda V0a \in ty\_2Einteger\_2Eint.(ap (c\_2Emin\_2E\_40 (ty\_2Einteger\_2Eint\_REP a)))$

Let  $c\_2Einteger\_2Eint\_mul : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_mul \in (((ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum) (ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum))^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)}) \quad (8)$$

Let  $c\_2Einteger\_2Eint\_eq : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_eq \in ((2^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)})^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)}) \quad (9)$$

Let  $c\_2Einteger\_2Eint\_ABS\_CLASS : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_ABS\_CLASS \in (ty\_2Einteger\_2Eint)^{(2^{(ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)})} \quad (10)$$

**Definition 12** We define  $c\_2Einteger\_2Eint\_ABS$  to be  $\lambda V0r \in (ty\_2Epair\_2Eprod ty\_2Enum\_2Enum ty\_2Enum\_2Enum)$

**Definition 13** We define  $c\_2Einteger\_2Eint\_mul$  to be  $\lambda V0T1 \in ty\_2Einteger\_2Eint.\lambda V1T2 \in ty\_2Einteger\_2Eint$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (11)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \quad (12)$$

Assume the following.

$$(\forall V0t \in 2.(((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee (p V0t)) \Leftrightarrow (p V0t)))))) \quad (13)$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in ty\_2Einteger\_2Eint. (\forall V1y \in ty\_2Einteger\_2Eint. \\
& (\forall V2z \in ty\_2Einteger\_2Eint. (((ap (ap c\_2Einteger\_2Eint\_mul \\
& V0x) V1y) = (ap (ap c\_2Einteger\_2Eint\_mul V0x) V2z)) \Leftrightarrow ((V0x = (ap \\
& c\_2Einteger\_2Eint\_of\_num c\_2Enum\_2E0)) \vee (V1y = V2z))))))
\end{aligned} \tag{14}$$

**Theorem 1**

$$\begin{aligned}
& (\forall V0x \in ty\_2Einteger\_2Eint. (\forall V1y \in ty\_2Einteger\_2Eint. \\
& (\forall V2z \in ty\_2Einteger\_2Eint. ((\neg (V0x = (ap c\_2Einteger\_2Eint\_of\_num \\
& c\_2Enum\_2E0))) \Rightarrow ((V1y = V2z) \Leftrightarrow ((ap (ap c\_2Einteger\_2Eint\_mul \\
& V0x) V1y) = (ap (ap c\_2Einteger\_2Eint\_mul V0x) V2z))))))
\end{aligned}$$