

thm_2Einteger_2EINT__LE__CALCULATE (TM- cRxtgRBRfasoYsubCZXimNVPN95PS82W6)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \tag{3}$$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \tag{4}$$

Definition 1 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p\ (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o\ (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_21$ to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2)))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x)$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a})))\ P)$

Definition 5 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint.(ap\ (c_2Emin_2E_40\ (ty_2Einteger_2Eint_REP_CLASS\ a)))$

Let $c_2Einteger_2Eint_lt : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \tag{5}$$

Definition 6 We define $c_2Einteger_2Eint_lt$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.\lambda V1T2 \in ty_2Einteger_2Eint$

Definition 7 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 8 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Definition 9 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 10 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E$

Definition 11 We define $c_2Einteger_2Eint_le$ to be $\lambda V0x \in ty_2Einteger_2Eint.\lambda V1y \in ty_2Einteger_2E$

Assume the following.

$$\begin{aligned} & (\forall V0x \in ty_2Einteger_2Eint.(\forall V1y \in ty_2Einteger_2Eint. \\ & ((p (ap (ap c_2Einteger_2Eint_le V0x) V1y)) \Leftrightarrow ((p (ap (ap c_2Einteger_2Eint_lt \\ & V0x) V1y)) \vee (V0x = V1y)))))) \end{aligned} \tag{6}$$

Theorem 1

$$\begin{aligned} & (\forall V0x \in ty_2Einteger_2Eint.(\forall V1y \in ty_2Einteger_2Eint. \\ & ((p (ap (ap c_2Einteger_2Eint_le V0x) V1y)) \Leftrightarrow ((p (ap (ap c_2Einteger_2Eint_lt \\ & V0x) V1y)) \vee (V0x = V1y)))))) \end{aligned}$$