

thm_2Einteger_2EINT__LTE__ANTSVM

(TMaxh7iZTD3LScVTLmsnuhXqDgYzRuEbomR)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \tag{3}$$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})_{ty_2Einteger_2Eint}) \tag{4}$$

Definition 2 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p)$ of type $\iota \Rightarrow \iota$).

Definition 3 We define $c_2Ebool_2E_21$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})))$

Definition 5 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint.(ap (c_2Emin_2E_40 (ty_2Einteger_2Eint_REP$

Let $c_2Einteger_2Eint_lt : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})_{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \tag{5}$$

Definition 6 We define $c_2Einteger_2Eint_lt$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.\lambda V1T2 \in ty_2Einteger_2Eint$

Definition 7 We define c_Ebool_2E to be $(ap (c_Ebool_2E_21) 2) (\lambda V0t \in 2.V0t)$.

Definition 8 We define $c_Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 9 We define $c_Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_Emin_2E_3D_3D_3E V0t) c_Ebool_2E$

Definition 10 We define $c_Einteger_2Eint_le$ to be $\lambda V0x \in ty_2Einteger_2Eint.\lambda V1y \in ty_2Einteger_2E$

Definition 11 We define $c_Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_Ebool_2E_21) 2) (\lambda V2t \in$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \wedge (p V1t2)) \Leftrightarrow ((p V1t2) \wedge (p V0t1)))))) \quad (6)$$

Assume the following.

$$(\forall V0x \in ty_2Einteger_2Eint.(\forall V1y \in ty_2Einteger_2Eint. (\neg((p (ap (ap c_Einteger_2Eint_lt V0x) V1y)) \wedge (p (ap (ap c_Einteger_2Eint_le V1y) V0x))))))) \quad (7)$$

Theorem 1

$$(\forall V0x \in ty_2Einteger_2Eint.(\forall V1y \in ty_2Einteger_2Eint. (\neg((p (ap (ap c_Einteger_2Eint_le V0x) V1y)) \wedge (p (ap (ap c_Einteger_2Eint_lt V1y) V0x)))))))$$