

thm_2Einteger_2EINT__LT__ADD (TM- RndK9XJj1nDhdsd9VCjWTv6nu6jQgoGbN)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_21$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define $c_2Ebool_2E_21$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 7 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \tag{4}$$

Let $c_2Einteger_2Eint_of_num : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_of_num \in (ty_2Einteger_2Eint^{ty_2Enum_2Enum}) \tag{5}$$

Assume the following.

$$(\forall V0x \in ty_2Einteger_2Eint.((ap (ap c_2Einteger_2Eint_add (ap c_2Einteger_2Eint_of_num c_2Enum_2E0)) V0x) = V0x)) \quad (14)$$

Assume the following.

$$(\forall V0w \in ty_2Einteger_2Eint.(\forall V1x \in ty_2Einteger_2Eint.(\forall V2y \in ty_2Einteger_2Eint.(\forall V3z \in ty_2Einteger_2Eint.(((p (ap (ap c_2Einteger_2Eint_lt V0w) V1x)) \wedge (p (ap (ap c_2Einteger_2Eint_lt V2y) V3z)))) \Rightarrow (p (ap (ap c_2Einteger_2Eint_lt (ap (ap c_2Einteger_2Eint_add V0w) V2y)) (ap (ap c_2Einteger_2Eint_add V1x) V3z)))))))))) \quad (15)$$

Theorem 1

$$(\forall V0x \in ty_2Einteger_2Eint.(\forall V1y \in ty_2Einteger_2Eint.(((p (ap (ap c_2Einteger_2Eint_lt (ap c_2Einteger_2Eint_of_num c_2Enum_2E0)) V0x)) \wedge (p (ap (ap c_2Einteger_2Eint_lt (ap c_2Einteger_2Eint_of_num c_2Enum_2E0)) V1y)))) \Rightarrow (p (ap (ap c_2Einteger_2Eint_lt (ap c_2Einteger_2Eint_of_num c_2Enum_2E0)) (ap (ap c_2Einteger_2Eint_add V0x) V1y))))))$$