

thm\_2Einteger\_2EINT\_\_LT\_\_GT  
(TMWVnPA8fydqoFZXEsW4SpRRQDH74oH9Nr8)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow P \Rightarrow Q)$  of type  $\iota$ .

**Definition 2** We define `c_2Emin_2E_3D` to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define `c_2Ebool_2E_2T` to be  $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define `c_2Ebool_2E_21` to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})))$

**Definition 5** We define `c_2Ebool_2E_2F` to be  $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$ .

Let `ty_2Enum_2Enum` :  $\iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let `ty_2Epair_2Eprod` :  $\iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let `ty_2Einteger_2Eint` :  $\iota$  be given. Assume the following.

$$nonempty\ ty\_2Einteger\_2Eint \tag{3}$$

Let `c_2Einteger_2Eint__REP__CLASS` :  $\iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)})ty\_2Einteger\_2Eint) \tag{4}$$

**Definition 6** We define `c_2Emin_2E_40` to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p))$  of type  $\iota \Rightarrow \iota$ .

**Definition 7** We define `c_2Einteger_2Eint__REP` to be  $\lambda V0a \in ty\_2Einteger\_2Eint.(ap (c_2Emin_2E_40 (ty\_2Einteger\_2Eint\_REP\_CLASS)))$

Let  $c\_2Einteger\_2Etint\_lt : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Etint\_lt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)})^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum)}) \quad (5)$$

**Definition 8** We define  $c\_2Einteger\_2Eint\_lt$  to be  $\lambda V0T1 \in ty\_2Einteger\_2Eint.\lambda V1T2 \in ty\_2Einteger\_2Eint.$

**Definition 9** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in 2.))$

**Definition 10** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E\_7E$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (7)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (9)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (10)$$

Assume the following.

$$(\forall V0x \in ty\_2Einteger\_2Eint.(\forall V1y \in ty\_2Einteger\_2Eint.(\neg((p\ (ap\ (ap\ c\_2Einteger\_2Eint\_lt\ V0x)\ V1y)) \wedge (p\ (ap\ (ap\ c\_2Einteger\_2Eint\_lt\ V1y)\ V0x)))))) \quad (11)$$

**Theorem 1**

$$(\forall V0x \in ty\_2Einteger\_2Eint.(\forall V1y \in ty\_2Einteger\_2Eint.((p\ (ap\ (ap\ c\_2Einteger\_2Eint\_lt\ V0x)\ V1y)) \Rightarrow (\neg(p\ (ap\ (ap\ c\_2Einteger\_2Eint\_lt\ V1y)\ V0x))))))$$